



ARAB REPUBLIC OF EGYPT  
Ministry of Education and  
Technical Education  
Central Administration of Book Affairs

Arts Section

# General Mathematics

second secondary grade

Student book

First term

## Authors

Mr. Kamal Yones Kabsha

Prof. Dr. Afaf Abo Elfotouh

Mr. Cerafiem Elias Skander

Mr. Magdy Abdelfatah Essafty

Mr. Ossama Gaber Abd-El-Hafez

Revised by

Mr. Mohamed Farouk Mohamed

Mr. Samir Mohamed Sedawy



2019 - 2020

غير مصرح بتداول هذا الكتاب خارج وزارة التربية والتعليم والتعليم الفني

**First edition**      **2017 / 2018**  
**Deposit number**   **10555 / 2015**  
**I.S.B.N**              **978 - 977 - 706 - 012 - 7**

# Introduction

---

## بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

We are pleased to introduce this book to show the philosophy on which the academic content has been prepared. This philosophy aims at:

- 1 Developing and integrating the knowledgeable unit in Math, combining the concepts and relating all the school mathematical curricula to each other.
- 2 Providing learners with the data, concepts, and plans to solve problems.
- 3 Consolidate the national criteria and the educational levels in Egypt through:
  - A) Determining what the learner should learn and why.
  - B) Determining the learning outcomes accurately. Outcomes have seriously focused on the following: learning Math remains an endless objective that the learners do their best to learn it all their lifetime. Learners should like to learn Math. Learners are to be able to work individually or in teamwork. Learners should be active, patient, assiduous and innovative. Learners should finally be able to communicate mathematically.
- 4 Suggesting new methodologies for teaching through (teacher guide).
- 5 Suggesting various activities that suit the content to help the learner choose the most proper activities for him/her.
- 6 Considering Math and the human contributions internationally and nationally and identifying the contributions of the achievements of Arab, Muslim and foreign scientists.

**In the light of what previously mentioned, the following details have been considered:**

- ★ This book contains three domains: algebra, relations and functions, calculus and trigonometry. The book has been divided into related and integrated units. Each unit has an introduction illustrating the learning outcomes, the unit planning guide, and the related key terms. In addition, the unit is divided into lessons where each lesson shows the objective of learning it through the title You will learn. Each lesson starts with the main idea of the lesson content. It is taken into consideration to introduce the content gradually from easy to hard. The lesson includes some activities, which relate Math to other school subjects and the practical life. These activities suit the students' different abilities, consider the individual differences throughout Discover the error to correct some common mistakes of the students, confirm the principle of working together and integrate with the topic. Furthermore, this book contains some issues related to the surrounding environment and how to deal with.
- ★ Each lesson contains examples starting gradually from easy to hard and containing various levels of thoughts accompanied with some exercises titled Try to solve. Each lesson ends in Exercises that contain various problems related to the concepts and skills that the students learned through the lesson.
- ★ Each unit ends in Unit summary containing the concepts and the instructions mentioned and General exams containing various problems related to the concepts and skills, which the student learned through the unit.
- ★ Each unit ends in an Accumulative test to measure some necessary skills to be gained to fulfill the learning outcome of the unit.
- ★ The book ends in General exams including some concepts and skills, which the student learned throughout the term.

**Last but not least. We wish we had done our best to accomplish this work for the benefits of our dear youngsters and our dearest Egypt.**

---

# Contents

---

## Unit 1 Real Functions and Graphing Curves

1 - 1 Real functions	3
1 - 2 Monotony of functions	12
1 - 3 Even and odd functions	17
1 - 4 Graphical representation of functions and geometric transformations.	24
1 - 5 Solving the absolute value equations and inequalities	39
Unit summary	46
Accumulative test	48

## Unit 2 Exponents, Logarithms and their Applications

2 - 1 Rational exponents	52
2 - 2 Exponential function and its graphical representation	60
2 - 3 Solving the exponential equations	64
2 - 4 Logarithmic function and its graphical representation	69
2 - 5 Some properties of logarithms	74
Unit summary	81
Accumulative test	83

# Contents

---

## Unit 3 Limits

3 - 1 Introduction to limits	86
3 - 2 Finding the limit of a function algebraically	92
3 - 3 Limit of a function at infinity	99
Unit summary	106
Accumulative test	107

## Unit 4 Trigonometry

4 - 1 Sine rule	110
4 - 2 Cosine rule	121
Unit summary	132
Accumulative test	133



# Unit 1

## Real Functions and Graphing Curves



### Unit introduction

Functions have various types and they have important applications in different life domains such as astronomy, medicine, economy, seismography, geology and demography. We use the functions to calculate the variables of the weather and to predict the expected weather conditions for a period of time in the future or to identify a malfunction position in the heart using the graphs which the electrocardiogram device record. Furthermore, functions are used to fulfill the best profit through studying the two functions of profit and cost or the effect of the age categories on census. Functions are also used in athletic medicine to identify the ideal weight (weight = length (cm) - 100) or to calculate the ratio of fats in the body.

In general, functions are commonly used in the industry to study the effect of the different variables on the production quality.

The Swiss scientist Leonard Euler (1707 - 1783) is considered one of the most prominent of the eighteenth century in mathematics and physics. He had been credited with using the symbol  $y = f(x)$  to express the function. He had considered that the function is a correlation between the elements of two sets with a relation that allows to calculate a variable value related to  $Y$  for another independent one. He had converted all the trigonometric ratios which ancient Egyptians, Babylonians and Arabs had excelled into trigonometric functions. In this unit, you are going to learn different forms of the real functions, their behaviour and their graphical representation using the geometrical transformations and graphical programs and to use the real functions in solving life and mathematical problems in different fields.



### Unit outcomes

By the end of the unit, the student should be able to:

- ✦ Identify the concept of the real function.
- ✦ Determine the domain, co-domain and range of the real functions.
- ✦ Deduce the monotony of the real functions of a real variable (increasing functions - decreasing functions - constant functions).
- ✦ Identify the type of the real function whether it is odd or even.
- ✦ identify polynomial functions.
- ✦ Graph the curves of (quadratic function - modulus functions - cubic function - rational function) and deduce the properties of each.
- ✦ Deduce the effect of the following transformations:  $f(x \pm a) \pm b$  and  $f(x \pm b) \pm c$  on the previous functions.
- ✦ Apply the previous transformation on graphing the curves of the real function.
- ✦ Solve equations in the form of:  $ax + b = c$ ,  $ax + b = |fx + c|$ .
- ✦ Solve inequalities in the form of:  $ax + b < c$  and  $ax + b \leq c$ ,  $ax + b \geq c$  and  $ax + b \leq c$ .
- ✦ Use the real functions to solve math and life problems in different fields.
- ✦ Relate what they learned about the effect of the previous transformations on the trigonometric functions in the form of activities.
- ✦ Investigate the graphical representation of the real function which have been previously learned and the effect of the previous transformation using the "Geogebra" program as cooperative work and activity.

## Key terms

- Real Function
- Domain
- Co-domain
- Range
- Vertical Line
- Piecewise-Defined Function
- Even Function
- Odd Function
- Monotony of Function
- Increasing Function
- Decreasing Function
- Constant Function
- polynomial Function
- Absolute Value Function
- Rational Function
- Asymptote
- Transformation
- Translation
- Reflection
- Stretching
- Graphical Solution

## Unit Lessons

- Lesson (1 - 1): Real functions.
- Lesson (1 - 2): Monotony of functions.
- Lesson (1 - 3): Even and odd functions.
- Lesson (1 - 4): Graphical representation of functions and geometric transformation.
- Lesson (1 - 5): Solving the equations and inequalities of absolute value.

## Materials

Computer set with graphic programs-  
graphic calculator-Scientific calculator

## Unit planning guide

### Real functions and graphing curves

#### Real function

Domain of function

Range of function

#### Monotony of function

Increasing function

Decreasing function

Constant function

#### Symmetry of function

Even function

Odd function

#### Graphical representation of functions and geometric transformation

#### Life applications

Solving equations

Solving inequations



## Real functions

## 1 - 1

**You will learn**

- ▶ The concept of the real function
- ▶ Vertical line test
- ▶ The piecewise defined function (defined with more than a rule).
- ▶ Identifying the domain and range of the real function.
- ▶ Operations on the functions.

**Key terms**

- ▶ Function
- ▶ Domain
- ▶ Co-domain
- ▶ Range
- ▶ Arrow diagram
- ▶ Cartesian diagram
- ▶ Vertical line
- ▶ Piecewise defined function
- ▶ Rule of the function.

**Materials**

- ▶ Scientific calculator
- ▶ Computer with graphic programs

**Explore**


You have previously learned the concept of the function and known that the function is a relation between two non-null sets  $X$  and  $Y$  so that each element of  $X$  is identified with one element of  $Y$  and denoted by one of these symbols  $f, g, h, \dots$

If we denote a function of set  $X$  to set  $Y$  with symbol  $f$ , it's mathematically written as :

$f: X \longrightarrow Y$  and read  $f$  function from  $X$  to  $Y$ . Notice:

- 1- For each element  $x \in X$ , one element of  $y \in Y$  is identified by the rule of the function  $f$  and written as:  $y = f(x)$

- 2- The set  $X$  is called the domain of the function and the set  $Y$  is called the co-domain of the function.



- 3- The set  $\{y = f(x) : x \in X\}$

is called the range of the function and known as the set of the image of the elements of the domain of the function.


Definition

**Real function:**

The function  $f$  is called a real function if each of its domain and co-domain are the set of the real numbers  $\mathbb{R}$  or a subset of it.


**Example**

- 1 The relation from set  $X$  to set  $Y$  represented in the arrow diagram opposite represents a function where: set  $X$  is the domain of the function =  $\{1, 2, 3, 4\}$  and the set  $Y$  is the co-domain of the function =  $\{5, 6, 7, 8, 9\}$ , whereas the set of elements  $\{6, 8, 9\}$  is known as the range of the function.

**Try to solve**

- 1 Which of the following relations shown by the illustrated arrow

diagrams represents a function and which does not represent a function? Write down the domain and range in case the relation represents a function:



### The graphical representation of the functions


If  $f: X \rightarrow Y$ , then the set of ordered pairs which satisfies the rule of the function is called the function. I.e. the function  $f = \{ (x, y) : x \in X, y \in Y \text{ and } y = f(x) \}$

By representing these ordered pairs on the cartesian diagram, we draw the graphical form of the function or the curve of the function.

In example (1) : The function  $f = \{ (1, 6), (2, 8), (3, 9), (4, 6) \}$ .

#### Notice:


- 1- The graphical form of the function is a set of separated points.
- 2- The vertical line passing at each element of the elements of the function domain intersects its graphical representation at an only point.




#### Learn

### The vertical line test

If the vertical line is found at each element of the domain elements, it passes through an only point of the points representing the relation, the relation was a function from  $X \rightarrow Y$



une fonction



n'est pas une fonction



#### Example

### Identifying the relation representing a function

- ② In each of the following figures, show whether  $y$  represents a function in  $x$  or not .




Figure (1)




Figure (2)




Figure (3)




Figure (4)



#### Solution

Figure (1) represents a function

Figure (2) does not represent a function because the vertical line passing through point (1, 0) intersects the graph at infinite number of points.

Figure (3) represents a function.

Figure (4) does not represent a function because there is a vertical line intersects the curve at more than a point.

**P Try to solve**

2 Show which of the following relations represents a function from  $X \rightarrow Y$ . Why?




Figure (1)




Figure (2)




Figure (3)





Figure (4)

**Example Identifying the range of the function.**


- 3 a If  $f: [1, 5] \rightarrow \mathbb{R}$  where  $f(x) = x + 1$   
Graph the function  $f$  and deduce the range of the function
- b If  $g: [1, 5[ \rightarrow \mathbb{R}$  where  $g(x) = x + 1$   
Graph the function  $f$  and deduce the range of the function.

**Solution**

- a Function  $f$  is a linear function whose domain is  $[1, 5]$ . It is represented by a line segment whose two ends are  $(1, f(1))$  and  $(5, f(5))$  i.e. the two points  $(1, 2)$  and  $(5, 6)$ .  
**The range of function  $f = [2, 6]$**   
It is the set of  $y$  coordinates for all the points which belong to the curve of the function.



- b The function  $g$  is a linear function whose domain is  $[1, 5[$ . It is clear that  $g(x) = f(x)$  for each  $x \in [1, 5[$  to be graphically represented by a line segment on a way that one of its ends is point  $(1, 2)$ , and the other end  $(5, 6)$  is to be ignored from the graph by placing an open circle at this point.  
The range of the function  $g = [2, 6[$



**Try to solve**

- 3 a If  $f: [1, \infty[ \rightarrow \mathbb{R}$ , where  $f(x) = 1 - x$   
Graph the function  $f$ , and deduce its range.
- b If  $g: ]-\infty, -1[ \rightarrow \mathbb{R}$ , where  $g(x) = 1 - x$   
Graph the function  $g$  and deduce its range.

**Piecewise - defined function****Work together**

To decrease the consumption of electricity, water and gas, the monthly consumption is calculated with respect to special categories relating the consumption amount to its value. The table opposite illustrates the prices of the monthly consumption categories for the natural gas at homes in piasters. Calculate the value of a home consumption of the natural gas in piasters with a classmate for the following quantities:

Monthly consumption in (m <sup>3</sup> )	Price per piasters
till 25	40
More than 25 till 50	100
More than 50	150

1- 30 cubic meters monthly.

2- 60 cubic meters monthly.

[Taxes and service are added after the monthly consumption is calculated]

**Notice:** The function  $f$  of calculating the value of the consumption  $x$  cubic meter of gas monthly where  $x \in \mathbb{R}$  can be written as follows:

$$f(x) = \begin{cases} 40x & \text{when } 0 \leq x \leq 25 \\ 100x - 1500 & \text{when } 25 < x \leq 50 \\ 150x - 4000 & \text{when } x > 50 \end{cases}$$

It is a **piecewise - defined function** (Defined by more than a rule)

**Learn**

The piecewise defined function is a real function where each subset of its domain has a different definition base.

**Try to solve**

- 4 Check your answer using the function above in **Work together**, then calculate the gas monthly consumption for the following quantities:
- a 15 cubic metres      b 40 cubic metres      c 54 cubic metres

**Graphing piecewise- defined function :****Example**

- 4 If  $f(x) = \begin{cases} 3 - x & \text{when } -2 \leq x < 2 \\ x & \text{when } 2 \leq x \leq 5 \end{cases}$


Graph and deduce the domain and range of the function.

**Solution**

The function  $f$  is defined on two intervals and  $f(x)$  is identified by two rules:

**First rule :**  $f_1(x) = 3 - x$  when  $-2 \leq x < 2$  i.e. on interval  $[-2, 2[$ .

It is up for a linear function represented by a line segment whose two ends are the two points  $(-2, 5)$  and  $(2, 1)$  by placing an open circle at point  $(2, 1)$  because  $2 \notin [-2, 2[$  as shown in the figure opposite.



**Second rule :**  $f_2(x) = x$  when  $2 \leq x \leq 5$  i.e. on interval  $[2, 5]$

It is up for a linear function represented by a line segment whose two ends are the two points  $(2, 2)$  and  $(5, 5)$  and the domain of the function  $f = [-2, 2[ \cup [2, 5] = [-2, 5]$

**From the graph, we deduce :**

the domain of the function  $f = [-2, 5]$





the range of the function  $f = ]1, 5]$

**Try to solve**


5 If  $f(x) = \begin{cases} x - 1 & \text{when } -2 \leq x < 0 \\ x + 1 & \text{when } x \geq 0 \end{cases}$

Graph the function and deduce its domain and range.

6 Deduce the domain and range of each function in the following graphs.



**Notice**



in the graph representing the function  $f$ , the domain of the function =  $[a, b]$  the range of the function =  $[c, d]$

### Identifying the domain of the real functions and the operations on them

The domain of the function is identified from its definition base or its graph.

#### Example Determining Domain of the function

5 Determine the domain of the following real functions defined by the next rules:

a  $f_1(x) = \frac{x+3}{x^2-9}$

b  $f_2(x) = \sqrt{x-3}$

c  $f_3(x) = \sqrt[3]{x-5}$

d  $f_4(x) = \frac{1}{\sqrt{3-x}}$

#### Solution

- a The function  $f_1$  is not defined when the denominator = 0, so we place  $x^2 - 9 = 0$  i.e.  $x = \pm 3$

Thus, the domain of the function  $f_1$  is  $\mathbb{R} - \{-3, 3\}$

- b The domain of the function  $f_2$  is all the values of  $x$  which make the value of the number inside the root positive or zero. I.e. the values of  $x$  which satisfy  $x - 3 \geq 0$

$\therefore x - 3 \geq 0 \quad \therefore x \geq 3$ ,  $\therefore$  the domain of the function  $f_2 = [3, \infty[$



- c  $f_3(x) = \sqrt[3]{x-6}$ , index of the root is an odd number and the domain of  $f_3 = \mathbb{R}$

- d  $f_4$  is defined when  $3 - x > 0$

Thus, the domain of  $f_4$  is  $]-\infty, 3[$



#### Notice:

If  $f(x) = \sqrt[n]{g(x)}$  where  $n \in \mathbb{Z}^+$  and  $n > 1$ , then  $g(x)$  is polynomial

**First:** When  $n$  is an odd number, the domain of the function  $f = \mathbb{R}$

**Second:** When  $n$  is an even number, the domain of the function  $f$  is the set of the values of  $x$  such that:  $g(x) \geq 0$

#### Try to solve

7 Determine the domain of the following real functions defined by the next rules:

a  $f_1(x) = \frac{2x+3}{x^2-3x+2}$

b  $f_2(x) = \sqrt{x-2}$

c  $f_3(x) = \sqrt[3]{x-5}$

d  $f_4(x) = \frac{5}{\sqrt{x+4}}$

**Critical thinking:** If the domain of the function  $f$  where  $f(x) = \frac{2}{x^2-6x+k}$  is  $\mathbb{R} - \{-3\}$ , find the value of  $k$ .

**Activity****Operations on functions**

If  $f_1$  and  $f_2$  are two functions whose two domains are  $m_1$  and  $m_2$  respectively, then:

- 1  $(f_1 \pm f_2)(x) = f_1(x) \pm f_2(x)$ , domain of  $(f_1 \pm f_2)$  is  $m_1 \cap m_2$
- 2  $(f_1 \cdot f_2)(x) = f_1(x) \cdot f_2(x)$ , domain of  $(f_1 \cdot f_2)$  is  $m_1 \cap m_2$
- 3  $(\frac{f_1}{f_2})(x) = \frac{f_1(x)}{f_2(x)}$  where  $f_2(x) \neq 0$  domain of  $(\frac{f_1}{f_2})$  is  $(m_1 \cap m_2) - Z(f_2)$   
where  $Z(f_2)$  is the set of zeros of  $f_2$

**We notice that:** in all the previous cases, the domain of the new function equals the intersection of the two domains of  $f_1$  and  $f_2$  except for the values which make  $f_2(x) = 0$  in the division operation.

- If**  $f_1 : \mathbb{R} \rightarrow \mathbb{R}$  where  $f_1(x) = 3x - 1$   
 $f_2 : [-2, 3] \rightarrow \mathbb{R}$  where  $f_2(x) = x - 3$

**First:** Find the rule and domain for each of the following functions:

- a**  $(f_1 + f_2)$       **b**  $(f_1 - f_2)$       **c**  $(f_1 \cdot f_2)$       **d**  $(\frac{f_1}{f_2})$


**Second:** Calculate the numerical value of each (if possible):

- a**  $(f_1 + f_2)(3)$       **b**  $(f_1 - f_2)(-3)$       **c**  $(f_1 \cdot f_2)(-2)$   
**d**  $(f_1 \cdot f_2)(2)$       **e**  $(\frac{f_1}{f_2})(4)$       **f**  $(\frac{f_1}{f_2})(-1)$


**Exercises 1 - 1**

**Choose the correct answer:**


- ① The relation shown by the following arrow diagrams which represents a function is :




- 2 The relation shown in the following graphical figures which does not represent a function is :




a



b



c







d

- 3 The relation shown by the set of the ordered pairs which does not represent a function is:

- a  $\{(1, 3), (3, 5), (5, 7), (7, 9)\}$       b  $\{(2, 3), (3, 4), (2, 1), (3, 5)\}$   
 c  $\{(0, 3), (1, 3), (2, 3), (3, 3)\}$       d  $\{(-3, 5), (-1, 5), (0, 5), (2, 5)\}$

Answer the following:

- 4 If  $f: X \rightarrow \mathbb{R}$  and  $X = \{1, 2, -2, -3\}$   
 Find the range of the function if  $f(x) = 5x - 3$
- 5 If  $g: \{1, 2, 3, 4, 5\} \rightarrow \mathbb{Z}^+$  where  $g(x) = 4x - 3$   
 a Write down the range of the function      b If  $g(k) = 17$ , find the value of  $k$
- 6 From the graph, deduce the domain and range of the function in each of the following:



- 7 Determine the domain of the function  $f$  where  $f(x) = \begin{cases} x - 1 & \text{when } 2 < x \leq 4 \\ -1 & \text{when } -2 \leq x \leq 2 \end{cases}$

Graph the function, then deduce its range.

- 8 Graph the function  $f$  where :

$$f(x) = \begin{cases} x + 3 & \text{when } x \geq 2 \\ 2x - 1 & \text{when } x < 2 \end{cases} \quad \text{then deduce its range.}$$

- 9 If  $f(x) = \begin{cases} 2x + 3 & \text{when } -2 \leq x < 0 \\ 1 - x & \text{when } 0 \leq x \leq 4 \end{cases}$

Graph the function  $f$  and deduce its range



$$10 \text{ If } f(x) = \begin{cases} x+1 & \text{when } -3 \leq x < 0 \\ x+2 & \text{when } 0 \leq x \leq 3 \end{cases}$$

Graph the function  $f$  and deduce its range.

$$11 \text{ If } f(x) = \begin{cases} -4x+3 & \text{when } x < 3 \\ -x^3 & \text{when } 3 \leq x \leq 8 \\ 3x^2+1 & \text{when } x > 8 \end{cases}$$

Find :

a)  $f(2)$

b)  $f(3)$

c)  $f(10)$

12 **Trade:** The function  $f$ , where :

$$f(x) = \begin{cases} \frac{5}{2}x & \text{When } 0 \leq x \leq 5000 \\ 2^x + 2500 & \text{When } 5000 < x \leq 15000 \\ \frac{3^x}{2} + 10000 & \text{When } 15000 < x \leq 60000 \end{cases}$$

represents the amount of money charged by a company to distribute an electrical appliance in L.E. If  $x$  represents the number of distributed appliances, find :

a)  $f(5000)$

b)  $f(10000)$

c)  $f(50000)$

13 **Geometry:** If  $S$  is the perimeter of a square whose side lengths is  $\ell$ , write the perimeter of the square as a function in its side length is ( $\ell$ ) then find :

a)  $S(3)$

b)  $S\left(\frac{15}{4}\right)$

14 **Geometry:** If  $M$  is the area of a circle whose radius lengths is  $r$ , Write the area of a circle as a function in its radius lengths  $r$   $M(r)$  then find  $M\left(\frac{1}{2}\right)$  and  $M(5)$ .

15 Determine the domain for each of the following real functions defined by the following rules:

a)  $f(x) = \frac{x+3}{x^2-5x+6}$

b)  $f(x) = \frac{x+1}{x^3+1}$

c)  $f(x) = \sqrt{x-2}$

d)  $f(x) = \sqrt{4-x^2}$


# 1 - 2 Monotony of Functions



## Think and discuss

The opposite graph shows the temperatures recorded in Cairo On a day. Observe the change of degrees over the time, then find using the graph:

- The periods when the temperature decreases.
- The periods when the temperature increases.
- The period when the temperature is constant. Characteristics of the curves help us know the behaviour of the function  $f$  and identify whether the function  $f(x)$  is increasing, decreasing or constant. It is called the monotony of the function.



## You will learn

- Monotony of functions.
- Using graphical program such as (GeoGebra) to graph the function curve.

## Key terms

- Monotony
- Increasing function
- Decreasing function
- Constant function



## Learn


The function  $f$  is

Said to be increasing on the interval  $]a, b[$

If each of  $x_1$  and  $x_2 \in ]a, b[$

where:  $x_2 > x_1$

then:  $f(x_2) > f(x_1)$




The function  $f$  is

Said to be decreasing on the interval  $]c, d[$

If each of  $x_1$  and  $x_2 \in ]c, d[$

where:  $x_2 > x_1$

then:  $f(x_2) < f(x_1)$




The function  $f$  is said to be constant on the interval  $]l, m[$

If each of:  $x_1$  and  $x_2 \in ]l, m[$

where  $x_2 > x_1$

then:  $f(x_2) = f(x_1)$



## Materials


- Scientific calculator
- Graphic programs

**Example**

- 1 Investigate the monotony of the function represented in the figure opposite.

**Solution**


- the function is decreasing in the interval  $]-\infty, 0[$
- the function is increasing in the interval  $]0, 2[$
- the function is constant in the interval  $]2, \infty[$



**Try to solve**




- 1 In the opposite figure:

Investigate the intervals in which the function is increasing, decreasing and constant.



**Example**

- 2 Each of the following graphs illustrate the curve of the function  $f: X \rightarrow Y$ . Deduce the domain and range of the function, then investigate its monotony






**Solution**

- a** The domain of  $f = \mathbb{R} = ]-\infty, \infty[$ , range of  $f = ]-\infty, \infty[$  the function increases in  $]-\infty, \infty[$
- b** The domain of  $f = ]-\infty, 2[ \cup ]2, +\infty[ = ]-\infty, \infty[$ , range of  $f = \mathbb{R}$   
the function increases in  $]-\infty, 2[$  and also in  $]2, \infty[$
- c** The domain of  $f = ]-\infty, 1[ \cup ]2, +\infty[$ , range of  $f = ]-\infty, 4[$   
the function is constant in  $]-\infty, 1[$ , and decreases in  $]2, \infty[$

**Try to solve**

- 2 Deduce the domain and range of the function, then investigate its monotony in each of the following graphs:



**Using the graphical programs to study the properties of the functions**


There are a lot of graphical programs to represent the functions graphically. The free GeoGebra for tablets or computers is one of the most famous.


**Activity**

Use the GeoGebra program to represent the function  $f$  graphically where:  $f(x) = x^3 - 3x + 2$ . Find the domain, range and investigate its monotony from the graph.


To do this activity, follow the next steps:


- 1- Open the algebraic window and the graph from GeoGebra, then press **Graphics** and choose  to reach the shown window in Fig (1).



- 2- In the algebraic window, write the rule of the function:  $f(x) = x^3 - 3x + 2$ , then inter (input) as follows:



Then press  the curve of the function appears in graphical window and the rule of the function in algebraic window as shown in Fig (2)



3- To determine points on the curve of a function, choose **A** from the tool bar then a new point from the menu. Move the pointer until it reaches the point determined on the curve. Press enter and the point will appear on the curve in the graphical window and the point coordinate appears in the algebraic window as shown in Fig (3).





Fig (3)

From the graph, we find :

- a) Domain of  $f = ]-\infty, \infty[$  and range of  $f = ]-\infty, \infty[$
- b) The function is increasing in  $]-\infty, -1[$ , decreasing in  $] -1, 1[$ , and increasing in  $]1, \infty[$ .



**Application**

Use GeoGebra to graph  $f(x) = 3x - x^3$ , then investigate the monotony of the function from the graph.

**Exercises 1 - 2**

1 Find the range and investigate the monotony of each of the following functions from the following graphs:




Fig (1)




Fig (2)




Fig (3)










Fig (4)

- 2 Determine the domain for each of the represented functions in the following graphs, then write the range and investigate its monotony:



- 3 If  $f: [-2, 6] \rightarrow \mathbb{R}$

$$f(x) = \begin{cases} 4 - x & \text{When } x < 1 \\ x & \text{When } 1 \leq x \leq 6 \end{cases}$$

Graph the function  $f$ , then deduce its range and investigate its monotony.

- 4 Use a graphical program to graph the curve of the function  $f$  in each of the following, then deduce its range and investigate its monotony.

a  $f(x) = x^2 - 5$

b  $f(x) = 4 - x^2$

c  $f(x) = (x - 1)^2 + 1$

d  $f(x) = x^3$

e  $f(x) = x^3 - 3x$

f  $f(x) = \frac{-1}{x - 2}$

# Even and Odd functions 1 - 3

## You will learn

- Symmetry in curves of functions.
- Even functions
- Odd functions

## Key terms

- Symmetry
- even function
- odd function


## Materials

- Scientific calculator
- Graphic programs


The graph of the function  $f$  where  $y = f(x)$  may be distinguished by geometrical characteristics that can be easily noticed from the graph. These characteristics can be used to study the functions and their applications. Symmetry around  $y$ -axis or around the origin point are of the most popular characteristics.

### Preface

You have previously learned the symmetry around a straight line where the figure can be folded around the straight line to make the two halves of the curve be congruent completely and you have also learned the symmetry around the origin point.



Symmetry around  $y$ -axis  
Figure (1)



Symmetry around the origin point.  
Figure (2)

### In figure (1):


The point  $(-x, y)$  which lies on the graph of the function curve is the image of point  $(x, y)$  which also lies on the same graph by reflection around  $y$ -axis.

### In figure (2):


The graph of the relation between  $x$  and  $y$  shows the symmetry of the curve around the origin point where point  $(-x, -y)$  is the image of point  $(x, y)$  which lies on the same curve.

### Try to solve


- 1 In the following figures, show which curve is symmetric around  $y$ -axis and which is symmetric around the origin point.



(a)



(b)



(c)

### Critical thinking:

Are curves of all functions symmetric around  $y$ -axis or around the origin point only? **Explain.**

## Even and Odd Functions



### Learn

**Even function :** It is said the function  $f: X \longrightarrow Y$  is an even function if  $f(-x) = f(x)$ , for each  $x, -x \in X$  and the curve of the even function is symmetric around  $y$ -axis.

**Odd function :** It is said the function  $f: X \longrightarrow Y$  is an odd function if  $f(-x) = -f(x)$ , for each  $x, -x \in X$  and the curve of the odd function is symmetric around the origin point.

**Notice :** A lot of functions are neither even nor odd.

When you investigate the type of the function whether it is even or odd, the condition of belonging the two elements  $x$  and  $-x$  to the domain of the function should be satisfied. If this condition is not satisfied, the function is neither even nor odd without finding  $f(-x)$ .



### Example

1 Investigate the type of the function  $f$  in each of the following and show whether it is odd or even.

a  $f(x) = x^2$

b  $f(x) = x^3$

c  $f(x) = \sqrt{x+3}$

d  $f(x) = \cos x$

### Solution

a  $f(x) = x^2$ , and the domain of  $f = \mathbb{R}$

for each  $x$  and  $-x \in \mathbb{R}$ , then  $f(-x) = (-x)^2 = x^2$

thus:  $f(-x) = f(x)$

$\therefore f$  is even function

b  $f(x) = x^3$ , and the domain of  $f = \mathbb{R}$

for each  $x$  and  $-x \in \mathbb{R}$ , then:  $f(-x) = (-x)^3 = -x^3$

thus:  $f(-x) = -f(x)$

$\therefore f$  is odd function

### Important remark :

The function  $f: \mathbb{R} \longrightarrow \mathbb{R}$  and  $f(x) = ax^n$  where  $a \neq 0$  and  $n \in \mathbb{Z}^+$  is called exponential function. The function is even when  $n$  is an even number and is odd when  $n$  is an odd number.



c  $f(x) = \sqrt{x+3}$ , and the domain of  $f = [-3, \infty[$

**Notice that**  $4 \in [-3, \infty[$  while  $-4 \notin [-3, \infty[$

$\therefore$  The function is neither even nor odd.

d  $f(x) = \cos x$ , and the domain of  $f = \mathbb{R}$

$\therefore$  for each  $-x$  and  $x \in \mathbb{R}$ , then  $f(-x) = \cos(-x) = \cos x$

**I.e.**  $f(-x) = f(x)$   $\therefore f$  is an even function



### Remember

$$\sin(-x) = -\sin x$$

$$\cos(-x) = \cos x$$

$$\tan(-x) = -\tan x$$

### Try to solve

2 Investigate the type of each function in the following functions and show whether it is even, odd or otherwise.

a  $f(x) = \sin x$

b  $f(x) = x^2 + \cos x$

c  $f(x) = x^3 - \sin x$

d  $f(x) = x^2 \cos x$

e  $f(x) = x^3 \sin x$

f  $f(x) = x^3 \cos x$

g  $f(x) = x^3 + x^2$

h  $f(x) = \sin x + \cos x$

i  $f(x) = \sin x \cos x$

**What do you infer ?**

### Important properties :

If each of  $f_1$  and  $f_2$  are even functions and each of  $g_1$  and  $g_2$  are odd functions, then :

1)  $f_1 + f_2$  is even function

2)  $g_1 + g_2$  is odd function

3)  $f_1 \times f_2$  is even function

4)  $g_1 \times g_2$  is even function

5)  $f_1 \times g_2$  is odd function




6)  $f_1 + g_2$  is neither even nor odd

Use these properties to verify your answers in **Try to solve**.



### Example

2 Each of the following graphs illustrates the curve of the function  $f$ . Determine whether the function is even, odd or otherwise, then check your answer algebraically.



### Solution

- a**  $f(x) = x^3 + x$ , from the graph of the function  $f$ , we notice that :  
 The domain of  $f = \mathbb{R}$  and the curve of the function is symmetric around the origin point.

I.e. the function is odd

$$\therefore \text{each of } x \in \mathbb{R} \text{ and } -x \in \mathbb{R} \quad \therefore f(-x) = (-x)^3 + (-x)$$

**By simplifying :**

$$f(-x) = -x^3 - x$$

**take off (-1) a common factor**

$$f(-x) = -(x^3 + x)$$

$$f(-x) = -f(x)$$

I.e. the function is odd.

- b**  $f(x) = 2 - x^2$ , from the graph of the function  $f$ , we notice that:  
 the domain of  $f = [-2, 2]$  and the curve of the function is symmetrical around y-axis . I.e.  
 the function is even

$$\therefore \text{each of } x \in [-2, 2] \text{ and } -x \in [-2, 2] \quad \therefore f(-x) = 2 - (-x)^2$$

**By simplifying**

$$f(-x) = 2 - x^2$$

$$f(-x) = f(x)$$

I.e. the function is even

- c**  $f(x) = x^2 - 4x$ , from the graph of the function  $f$ , we notice that :  
 the domain of  $f = \mathbb{R}$  and the curve of the function is neither symmetric around y-axis nor  
 around the origin point. I.e. the function is neither even nor odd.

$x \in \mathbb{R}$ :

$$\therefore \text{Each of } x \in \mathbb{R} \text{ and } -x \in \mathbb{R} \quad \therefore f(-x) = (-x)^2 - 4(-x)$$

**By simplifying**

$$f(-x) = x^2 + 4x \neq f(x) \quad \therefore f \text{ is not an even function}$$

**But**

$$-f(x) = -x^2 + 4x$$


**Then**


$$f(-x) \neq -f(x) \quad \therefore f \text{ is not an odd function}$$

I.e. the function is neither even nor odd.

**P Try to solve**

- 3** Tell whether each of the functions represented in the following figures is even, odd or otherwise.






### Example

- 3 The opposite figure represents the curve of the function  $f$  where:

$$f(x) = \begin{cases} -\frac{1}{x} & \text{when } x < 0 \\ \frac{1}{x} & \text{when } x > 0 \end{cases}$$

Show that this function is even .



### Solution

From the graph, the curve is symmetric around  $y$ -axis. I.e. the function is even.

### Try to solve

- 4 Represent the function  $f$  where  $f(x) = \begin{cases} x + 2 & \text{when } x \geq -2 \\ -x - 2 & \text{when } x < -2 \end{cases}$  graphically.

then show whether the function is even, odd or otherwise.

## Exercises 1 - 3

- 1 Tell whether the symmetry of the curve is around  $x$ -axis,  $y$ -axis or origin point, then explain.




Fig (1)




Fig (2)




Fig (3)




Fig (4)




Fig (5)




Fig (6)

- 2 Find the range of each function and tell whether it is even, odd or otherwise.




Fig (a)




Fig (b)




Fig (c)




Fig (d)




Fig (e)




Fig (f)

- 3 Tell whether the function  $f$  is even, odd or otherwise.

a  $f(x) = x^4 + x^2 - 1$

b  $f(x) = 3x - 4x^3$

c  $f(x) = x^3 - \frac{1}{x}$

d  $f(x) = x^2 - 3x$

e  $f(x) = \frac{x^3 + 2}{x - 3}$

f  $f(x) = x \cos x$

- 4 If  $f_1, f_2$  and  $f_3$  are three real functions where  $f_1(x) = x^5, f_2(x) = \sin x$  and  $f_3(x) = 5x^2$ , tell which of the following functions is even, odd or otherwise.

a  $f_1 + f_2$

b  $f_1 + f_3$

c  $f_1 \times f_2$

d  $f_3 \times f_2$

- 5 Graph the curves for each of the defined functions, then tell whether it is even, odd or otherwise and investigate its monotony.

$$\text{a } f(x) = \begin{cases} 2 & \text{when } x > 0 \\ -2 & \text{when } x < 0 \end{cases}$$

$$\text{b } f(x) = \begin{cases} -x & \text{when } x \geq 0 \\ x & \text{when } x < 0 \end{cases}$$

$$\text{c } f(x) = \begin{cases} x - 1 & \text{when } x \geq 0 \\ 7x & \text{when } x < 0 \end{cases}$$

$$\text{d } f(x) = \begin{cases} x + 1 & \text{when } x \geq 0 \\ 1 - x & \text{when } x < 0 \end{cases}$$

- 6 Answer the following using the next graphs :




Fig (1)



Fig (2)




Fig (3)




Fig (4)

**First:** Complete the graph in figures (1) and (3) in your notebook to get an even function on its domain.

**Second:** Complete the graph in figures(2) and (4) in your notebook to get an odd function on its domain.

**Third:** Determine the domain and range of the function in each case , then investigate its monotony

**Polynomial functions**

You have previously learned the polynomial function whose base is in the form:  $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots + a_nx^n$

**where:**  $a_0, a_1, a_2, a_3, \dots, a_n \in \mathbb{R}, a_n \neq 0$  and  $n \in \mathbb{N}$

You knew that the domain and co-domain are the set of the real numbers  $\mathbb{R}$  (or a subset of it). As a result, these functions are called polynomial functions of  $n$  degree. The degree of a polynomial is the highest power of the independent variable  $x$ .

**Notice:**

- 1- If  $f(x) = a_0$  and  $a_0 \neq 0$  then  $f$  is called a constant polynomial function.
- 2- The polynomial functions of the first degree are called linear functions, of the second degree are called quadratic functions and of the third degree are called cubic functions.
- 3- Adding or subtracting functions of different powers and constants, we get a polynomial function.
- 4- Zeros of the polynomial function are the  $x$ -coordinates of the intersecting points of its curve with  $x$ -axis.

**Graphing the curves of the functions.****First: Polynomial functions:****Learn**


Here is the graphical representation of some polynomial functions :

- 1) **The simplest form of the linear function is:**

$$f(x) = x$$

it is a function  $f$  that joins the number itself and a straight line passing through point  $(0, 0)$ , and its slope = 1 represents it.

(**Check:** the range of  $f = \mathbb{R}$ ,  $f$  is odd and  $f$  is increasing in  $\mathbb{R}$ ).

**You will learn**

- The polynomial functions (linear function - quadratic function and cubic function)
- The absolute value function.
- The rational function
- using the geometrical transformation of the function  $f$  to graph the curves
  - $y = f(x) + a$
  - $y = f(x + a)$
  - $y = f(x + a) + b$
  - $y = -f(x)$
  - $y = af(x)$
  - $y = af(x + b) + c$
- Transformation of some trigonometric functions.

**Key terms**

- Transformation
- Translation
- Reflection
- Vertical
- Horizontal
- Asymptote

**Materials**


- Scientific calculator.
- Computer
- Graph program

- 2) The simplest form of the quadratic function  $f$  is :

$$f(x) = x^2$$

the quadratic function joins the number with its square. It is represented by an upward open curve, symmetrical around y-axis and the vertex point of the curve is  $(0, 0)$ .

(Check: the range of  $f = \mathbb{R}$ ,  $f$  is even,  $f$  is decreasing at  $]-\infty, 0[$  and increasing at  $]0, +\infty [$  )




- 3) The simplest form of the cubic function  $f$  is:

$$f(x) = x^3$$

The cubic function joins the number with its cubic.

It is represented by a curve whose symmetrical point is  $(0, 0)$ .

(Check: the range of  $f = \mathbb{R}$ ,  $f$  is odd and  $f$  is increasing in  $\mathbb{R}$ )



### Example

- 4) Graph the function  $f$  where:

$$f(x) = \begin{cases} x^2 & \text{when } x < 2 \\ 4 & \text{when } x > 2 \end{cases}$$

### Solution

- 1) When  $x < 2$  and  $f(x) = x^2$

We graph  $f(x) = x^2$  for each  $x \in ]-\infty, 2[$

Place an **open circle** at point  $(2, 4)$  as shown in **fig (1)**




Fig (1)

- 2) when  $x > 2$  and  $f(x) = 4$

We graph the constant function  $f(x) = 4$  for each  $x \in ]2, \infty [$  on the same graph **fig (2)**

Notice that the domain of the function  $f = \mathbb{R} - \{2\}$  and the range of  $f = ]0, \infty [$




Fig (2)

**P Try to solve**

- ① Graph the function  $f$  where:

$$f(x) = \begin{cases} x^2 & \text{when } x < 0 \\ x & \text{when } x \geq 0 \end{cases} \quad \text{then deduce the range of the function and investigate its monotony.}$$



**Learn**

**The Absolute Value Function**

the simplest form of the absolute value function is:

$$f(x) = |x|, x \in \mathbb{R}$$

It is defined as:


$$f(x) = \begin{cases} x & \text{when } x \geq 0 \\ -x & \text{when } x < 0 \end{cases}$$

**Notice:**  $|-3| = |3| = 3$ ,  $|0| = 0$ ,  $\sqrt{(-2)^2} = \sqrt{2^2} = 2$

**i.e.:**  $|x| \geq 0$ ,  $|-x| = |x|$ ,  $\sqrt{x^2} = |x|$

The function  $f$  is represented by two rays starting at point  $(0, 0)$ , the slope of the first = 1 and the slope of the other = -1.

(**Check:** the range of  $f = [0, \infty[$ ,  $f$  is even,  $f$  is decreasing in  $]-\infty, 0[$  and is increasing in  $]0, \infty[$ )



**Learn**

**Rational Function**


the simplest form of the rational function is:

$$f(x) = \frac{1}{x}, x \in \mathbb{R} - \{0\}$$

the function  $f$  joins the number with its multiplicative inverse. It is represented by a curve whose symmetrical point is  $(0, 0)$ . The curve consists of two parts one of them lies on first quadrant and the other lies on third quadrant and each part approaches the two axes and does not intersect them

( $x = 0$  and  $y = 0$  are two approaching lines of the curve)

(**Check:** The range of  $f = \mathbb{R} - \{0\}$ ,  $f$  is odd,  $f$  is decreasing in  $]-\infty, 0[$  and is decreasing in  $]0, \infty[$ )



**P Try to solve**

- ② Graph the function  $f$  where  $f(x) = \begin{cases} |x| & \text{when } x \leq 0 \\ \frac{1}{x} & \text{when } x > 0 \end{cases}$

From the graph, find the range and investigate the monotony of the function.



## Geometrical transformations of the function curves

### First : vertical translation of the function curve



#### Work together

#### Work with your classmate

- 1) Graph the curve of the function  $f: f(x) = x^2$  using Geogebra.
- 2) Place the pointer on the curve vertex and drag it vertically upward for one unit. Notice the change of the function rule to express a new function whose rule is:  
 $f(x) = x^2 + 1$  as shown in **figure (1)**.
- 3) Drag the curve vertex of the function to the points  $(0, 2)$  and  $(0, 3)$  then record your observations each time.
- 4) Drag the curve of  $f(x) = x^2$  vertically downward for two units and notice the change of the function rule to express a new function whose rule is :  $f(x) = x^2 - 2$  as shown in **Figure (2)**

**Think :** Show how  $f(x) = x^2 - 5$  can be graphed using the curve of  $f(x) = x^2$ ?

**Of the previous, we can notice that :**

If  $f(x) = x^2$ ,  $g(x) = x^2 + 1$  and  $h(x) = x^2 - 2$ , then :

- 1) The curve of  $g(x)$  is the same curve of  $f(x)$  by translation of a magnitude of one unit in the positive direction of  $y$ -axis.
- 2) The curve of  $h(x)$  is the same curve of  $f(x)$  by translation of a magnitude of two units in the negative direction of  $y$ -axis.

**Critical thinking :** Use the curve of  $f(x) = x^3$  to show how the following curves can be graphed:

**a**  $g(x) = x^3 + 4$

**b**  $h(x) = x^3 - 5$




Figure (1)




Figure (2)




Figure (3)



#### Learn

#### drawing the curve of $y = f(x) + a$

For any function  $f$ , the curve of  $y = f(x) + a$  is the same curve of  $y = f(x)$  by translation of a magnitude of  $a$  unit in the direction of  $\vec{oy}$  when  $a > 0$  and in the direction of  $-\vec{oy}$  when  $a < 0$ .

**Example**

- 5 The opposite figure shows the curves of the functions  $f$ ,  $g$  and  $h$  where each of  $g$  and  $h$  are an image of the function  $f$  by a vertical translation. Write the rules of  $g$  and  $h$ .


**Solution**

$\therefore$  The curve of the function  $g$  is the same curve of the function  $f$  by translation of a magnitude of three units in the direction of  $\vec{oy}'$




$$g(x) = f(x) - 3 \quad \therefore f(x) = |x| \quad \therefore g(x) = |x| - 3$$

$\therefore$  the curve of the function  $h$  is the same curve of the function  $f$  by translation of a magnitude of two units in the direction of  $\vec{oy}$

$$\therefore h(x) = f(x) + 2 \quad \therefore f(x) = |x| \quad \therefore h(x) = |x| + 2$$


**Try to solve**

- 3 The given figures show the curves of the functions  $f$ ,  $g$  and  $h$  where each of  $g$  and  $h$  are the image of the function  $f$  by a vertical translation. Write the rules of  $g$  and  $h$  in each figure.


**Second : Horizontal translation of the function curve:**
**Work together**
**Work with your classmate:**

- 1) Graph the function  $f: f(x) = |x|$  using GeoGebra by writing the rule of the function in the enter box as follows:  $\text{abs}(x)$  and press enter then the curve of the function will appear in the graphical window and its rule is  $f(x) = |x|$  in the algebraic window as in Fig (1)






Figure (1)



**Solution**


➤ The curve of  $g(x) = (x - 2)^2$  is the same curve of  $f(x) = x^2$  by translation of 2 units in the positive direction of  $x$ -axis and the vertex point of the curve is  $(2, 0)$ .



➤ The curve of  $h(x) = (x + 3)^2$  is the same curve of  $f(x) = x^2$  by translation of 3 units in the negative direction of  $x$ -axis and the vertex point of the curve is  $(-3, 0)$ .



**Try to solve**

- 4 Use the curve of the function  $f$  where  $f(x) = x^2$  to represent each of the two functions  $g$  and  $h$  where :

**a**  $g(x) = (x + 4)^2$

**b**  $h(x) = (x - 3)^2$

- 5 Write down the rule of the function  $f$  represented graphically in the following graphs :



**Critical thinking:** If  $f(x) = x^2$ , show how the curve of the function  $g$  where  $g(x) = (x - 3)^2 + 2$  can be graphed.

**Graphing the curve of  $y = f(x + a) + b$** 

**Of the previous, we deduce that:** The curve of  $y = f(x + a) + b$  is the same curve of  $y = f(x)$  by horizontal translation of a magnitude of  $a$  of units ( in the direction of  $\overrightarrow{ox'}$  when  $a < 0$  and in the direction of  $\overrightarrow{ox'}$  when  $a > 0$  ), then vertical translation of a magnitude of  $b$  of units ( In the direction of  $\overrightarrow{oy'}$  when  $b > 0$  and in the direction of  $\overrightarrow{oy'}$  when  $b < 0$  ).

**P Try to solve**

- 6 Use the curve of the function  $f$  where  $f(x) = x^2$  to represent each of the two functions  $g$  and  $h$  where :
- a  $g(x) = (x + 2)^2 - 4$                       b  $h(x) = (3 - x)^2 - 1$

**Example****Applying the geometrical transformations on graphing the function curves**


- 7 Graph the curve of the function  $g$  where  $g(x) = \frac{1}{x-1} + 3$ , then determine the range of the function and investigate its monotony from the graph

**Solution**

The curve of the function  $g$  is the same curve of the function  $f$  where  $f(x) = \frac{1}{x}$  by translation of a magnitude of one unit in the direction of  $\overrightarrow{OX}$  ( $a = -1 < 0$ ), then translation of a magnitude of three units in the direction of  $\overrightarrow{OY}$  and the point of symmetry of the function curve  $g$  is the point  $(1, 3)$  and the range of  $g = \mathbb{R} - \{3\}$

Monotony of the function  $g$ :




$g$  is decreasing in  $]-\infty, 1[$  and also decreasing in  $]1, \infty[$



**Critical thinking:** Can it be said that  $f(x) = \frac{1}{x-2} + 3$  is decreasing on its domain? Explain.


**P Try to solve**

- 7 Use the curve of the function  $f$  where  $f(x) = \frac{1}{x}$  and  $x \neq 0$  to represent each of the following:
- a  $g(x) = \frac{1}{x+2} + 1$                       b  $h(x) = \frac{2x-3}{x-2}$
- 8 Write down the rule of the function represented graphically by the following graphs:



**Remark:** The curve of  $f(x) = x^2 + 4x + 1$  can be graphed using the vertical and horizontal translation of the curve  $g(x) = x^2$  as follows.

$$\begin{aligned} f(x) &= x^2 + 4x + 1 \text{ by completing the square} \\ &= (x^2 + 4x + 4) - 3 \\ &= (x + 2)^2 - 3 \end{aligned}$$




i.e. the curve of the (given) function  $f$  is the same curve of the function  $g$  where  $g(x) = x^2$  by translation of a magnitude of two units in the direction of  $\overrightarrow{ox'}$ , then three units in the direction of  $\overrightarrow{oy'}$ . The opposite figure shows that.

**Application:** Graph the curve of the function  $f(x) = x^2 + 6x + 7$  using the vertical and horizontal translation of the function  $g(x) = x^2$ , then investigate the monotony of the function  $f$ .

### Third: The reflection of the function curve on x-axis

The following graphs illustrate the reflection of the curves of some standard functions on x-axis.



What do you notice? What do you infer?



### Learn

Graph the curve of  $y = -f(x)$

for any function  $f$ , the curve of  $y = -f(x)$  is the same curve of  $y = f(x)$  by reflection on x-axis.



### Example

#### Applying the geometrical transformations on graphing the curves

8 Use the curves of the standard functions to graph the functions  $g$ ,  $h$  and  $z$  where:

a  $g(x) = -(x - 3)^2$


b  $h(x) = 4 - |x+3|$

c  $z(x) = 2 - \frac{1}{x-3}$




### Solution


a The curve of  $g(x)$  is the reflection of the curve of  $f(x) = x^2$  on x-axis, then horizontal translation of a magnitude of three units in the direction of  $\overrightarrow{ox'}$ , and the vertex point of the curve is  $(3, 0)$  and the curve is open downward.



- b** The curve of  $h(x)$  is a reflection of the curve of  $f(x) = |x|$  on  $x$ -axis, then horizontal translation of a magnitude of 3 units in the direction of  $\vec{ox'}$  and vertical translation of a magnitude of four units in the direction of  $\vec{oy}$ . The starting point of the two rays is point  $(-3, 4)$  and the curve is open downward.



- c** The curve of  $z(x)$  is a reflection of the curve of  $f(x) = \frac{1}{x}$  on  $x$ -axis, then horizontal translation of a magnitude of three units in the direction of  $\vec{ox}$  and vertical translation of a magnitude of two units in the direction of  $\vec{oy}$  and the symmetrical point of the curve is  $(3, 2)$ .



**P Try to solve**

- 9** Graph the curve of the function  $g$  in each of the following where:

**a**  $g(x) = 3 - (x + 1)^2$

**b**  $g(x) = -(x - 3)^3$

**c**  $g(x) = 3 - |x - 5|$

Check your graph using a graphical program or the graphical calculator.

#### Fourth: Expanding the curve of the function



#### Work together

Graphing the curve of  $g(x) = a f(x)$

work with your classmate.

- 1)** Graph the curve of the function  $f : f(x) = x^2$  using GeoGebra and type the function rule in the input box as follows:



A new window will appear (Figure 1)

Select **Create sliders**

- 2)** Use the pointer of a values to select new values where  $1 < a$  and notice the motion of the function curve with respect to the curve of the function  $f$  for each  $x \in \mathbb{R}$  as in figure (2) and when  $1 > a$  as in figure (3)  
What do you notice ? what do you infer ?




Figure (1)




Figure (2)



## Learn

Graph the curve of  $y = a f(x)$ 

For any function  $f$ , the curve of  $y = a f(x)$  is a vertical stretch of the curve of  $y = f(x)$  if  $a > 1$  and vertical shrinking of the curve of  $y = f(x)$  if  $0 < a < 1$ .



## Example

- 9 Use the curve of the function  $f$  where  $f(x) = |x|$  to represent each of the two functions  $g$  and  $z$ .

a  $g(x) = 2|x|$

b  $z(x) = 2|x - 7| + 2$



## Solution

- a The curve of  $g(x)$  is a vertical stretch of the curve of the function  $f$  whose factor is  $a = 2 > 0$ . Thus, for each  $(x, y) \in f$  and  $(x, 2y) \in g$

- b The curve of  $z(x)$  is the same curve of  $g(x)$  by horizontal translation of a magnitude of 7 units in the direction of  $\overrightarrow{ox}$  and vertical translation of a magnitude of 2 units in the direction of  $\overrightarrow{oy}$ .





Fig (3)



## Try to solve

- 10 Use the curve of the function  $f$  where  $f(x) = x^2$  to represent the two functions  $g$  and  $z$ :

a  $g(x) = -\frac{1}{2}x^2$

b  $z(x) = 2 - \frac{1}{2}(x - 5)^2$

Check your graph using a graphical program or calculator, then determine the range of the function  $z$  and investigate its monotony.



## Activity

- Applying the geometrical transformations which you have learned in the previous algebraic functions on the sine and cosine functions.

## Trigonometric functions

## First: translation in the direction of x-axis:

- 1) Use GeoGebra and prepare programs in a way that the graduation on x-axis is in radian. this can be done by doing a (right click) on the mouse and choose the last line of x-axis and choose the graduation system ( $\pi$ ).
- 2) At the bottom of the program, type the order  $\sin(x)$  then press (enter) then you get the shape of the red curve.



You can control the color and thickness of the curve by pressing the mouse (left click), the color, thickness and the type of the line-dotted, slash or connected will appear upward the window.


3) Use the same way and type the order  $\sin(x + (3/\pi))$  i.e.:  $y = \sin(x + \frac{\pi}{3})$ , then press (enter) and color the curve in a different color.

4) Compare the two curves. What do you notice?

**From the graph, we deduce that:**

The curve of the sine function is translated horizontally to the left side on x-axis with a magnitude equals  $\frac{\pi}{3}$  (as in real function). we notice that the range of the constant function

$[-1, 1]$  is the same range of the function  $\sin x$ . We also notice that the function  $\sin(x + \frac{\pi}{3})$  is neither even nor odd because there is not symmetry to its curve around y-axis or origin point.



**Think:** What do you expect about the direction of the x-translation if the rule of the function is  $\sin(x - \frac{\pi}{3})$ ?

**Second: translation in the direction of y-axis**

1) Graph the curve of the function  $f$  where  $y = \sin x$ .


2) Graph the curve of the function  $g(x)$  where  $g(x) = \sin x + 2$

**in different color, then compare the shape of the two curves.**

**What do you notice?**



From the graph, we infer that:

The curve of the second function is the same curve of the function  $\sin(x)$  after translating it in a magnitude of 2 units upward. We notice that the range of the second function is  $[1, 3]$  because it is translated in a magnitude of 2 units in the positive direction of y-axis from the first function and the function  $\sin x + 2$  is neither even nor odd.



**Critical thinking:**

In each of the following graphs: describe the geometric transformations of the curve of the function  $f$  which graphs the curve of the function  $g$ , then write the rule of the function  $g$ , determine its range and investigate its monotony.





## Exercises 1 - 4



- 1 Graph the curve of the function  $f$ , then determine its range and investigate its monotony

a  $f(x) = \begin{cases} |x| & \text{when } x \leq 0 \\ x^2 & \text{when } x > 0 \end{cases}$

b  $f(x) = \begin{cases} 4 & \text{when } x < -2 \\ x^2 & \text{when } x \geq -2 \end{cases}$

c  $f(x) = \begin{cases} x^3 & \text{when } x < 1 \\ 1 & \text{when } x > 1 \end{cases}$

d  $f(x) = \begin{cases} \frac{1}{x} & \text{when } x < 0 \\ |x| & \text{when } x > 0 \end{cases}$

Choose the correct answer:

- 2 The curve of  $g(x) = x^2 + 4$  is the same curve of  $f(x) = x^2$  by translation of a magnitude of 4 units in the direction of :

a  $\vec{ox}$

b  $\vec{ox}'$

c  $\vec{oy}$

d  $\vec{oy}'$

- 3 The curve of the function  $g(x) = |x + 3|$  is the same curve of  $f(x) = |x|$  by translation of a magnitude of 3 units in the direction of:

a  $\vec{ox}$

b  $\vec{ox}'$

c  $\vec{oy}$

d  $\vec{oy}'$

- 4 The curve vertex point  $f(x) = (2 - x)^2 + 3$  is :

a (2, 3)

b (2, -3)

c (-2, 3)

d (-2, -3)

- 5 The symmetry point of the curve of the function  $f(x) = 2 - (x + 1)^3$  is :

a (1, 2)

b (-1, 2)

c (2, 1)

d (2, -1)

- 6 The symmetry point of the curve of the function  $f$  where  $f(x) = \frac{1}{x-3} + 4$  is :

a (3, -4)

b (-3, -4)

c (3, 4)


d (-3, 4)

Answer the following:


- 7 Graph the curve of the function  $f$  where  $f(x) = x^2$  then translate in the direction of the two coordinate axes  $x$  and  $y$  as shown in the opposite figure.

Write down the rule of the following functions:

➤  $g$ ,  $h$  and  $z$ .




- 8 In the figure opposite : Graph the curve of the function  $f$  where  $f(x) = x^3$ , then translate in the direction of the two coordinate axes  $x$  and  $y$ .  
Write down the rule of each of the following functions:



➤  $g, h$  and  $z$

- 9 In the figure opposite : Graph the curve of the function  $f$  where  $f(x) = |x|$  then translate in the direction of the two coordinate axes  $x$  and  $y$ .  
Write down the rule of each of the following functions :




➤  $g, h$  and  $z$


- 10 Graph the curve of the function  $f$  where  $f(x) = \frac{1}{x}$ , then translate in the direction of the two coordinate axes  $x$  and  $y$ .

Write down the rule of each function which the following curves represent:


a



b



c



- 11 Use the curve of the function  $f$  where  $f(x) = x^2$  to represent the following graphically:

a  $f_1(x) = x^2 - 4$

b  $f_2(x) = (x - 3)^2$

c  $f_3(x) = (x - 1)^2 - 2$

- 12 Use the curve of the function  $f$  where  $f(x) = |x|$  to represent the following graphically:

a  $f_1(x) = |x| + 1$

b  $f_2(x) = |x + 2|$

c  $f_3(x) = |x - 3| - 2$


➤ Then find the coordinates of the intersecting points of the curves with the two coordinate axes.

- 13 Use the curve of the function  $f$  where  $f(x) = x^3$  to represent the following graphically :
- a  $f_1(x) = f(x) - 3$       b  $f_2(x) = f(x - 2)$       c  $f_3(x) = f(x + 3) + 2$
- Then determine the symmetry point of the curve of each function.
- 14 If the function  $f$  where  $f(x) = \frac{1}{x}$  graph the function  $h$  and determine the symmetry point of the function curve :
- a  $g(x) = f(x - 3)$       b  $g(x) = f(x) + 2$       c  $g(x) = f(x - 2) + 2$
- 15 Use the curve of the function  $f$  where  $f(x) = x^2$  to represent the following graphically :
- a  $f_1(x) = 4 - x^2$       b  $f_2(x) = -(x - 3)^2$       c  $f_3(x) = 2 - (x + 3)^2$
- 16 use the curve of the function  $f$  where  $f(x) = |x|$  to represent the following graphically.
- a  $f_1(x) = 2 - |x|$       b  $f_2(x) = -|x + 5|$       c  $f_3(x) = 4 - |x - 2|$
- d  $f_4(x) = 2|x|$       e  $f_5(x) = -2|x - 1|$       f  $f_6(x) = 5 - 2|x + 2|$
- 17 Graph the curve of the function  $f$  in each of the following using the proper transformations, then investigate its monotony.

a  $f_1(x) = \begin{cases} x^2 + 2 & \text{when } x \geq 0 \\ -x^2 - 2 & \text{when } x < 0 \end{cases}$       b  $f_2(x) = \begin{cases} x^2 + 1 & \text{when } -4 \leq x < 0 \\ -x^2 - 1 & \text{when } 0 \leq x \leq 4 \end{cases}$

- 18 **Industry:** An iron gate whose two sides are 3 meters high and its arc is in the form of a part of the curve of the function  $f: f(x) = a(x - 2)^2 + 4$  has been designed as shown in the opposite figure, find:

- a Value of  $a$       b Maximal height of the gate  
c Width of the gate



- 19 **Trade:** A dealer pays 50 L.E for each ton getting in or out of his warehouse for loading or unloading the goods, write down the function representing the cost of loading or unloading, then represent it graphically.
- 20 **Urban communities:** rectangle-like pieces of land are specialized for youth housing in a new urban community. If the length of each is  $x$  meter and the area is  $400 \text{ m}^2$ .
- a Write down the rule of the function  $f$  which shows the width of the piece of land in terms of its length, then represent it graphically.
- b From the graph, find the width of the piece of land whose length is 25 meters, then check algebraically.

# Solving absolute value equations and inequalities

## 1 - 5

### You will learn

- ▶ Solve the modulus equations graphically.
- ▶ Solve the modulus equations algebraically.
- ▶ Solve the modulus inequalities graphically.
- ▶ Solve the modulus inequalities algebraically.
- ▶ Model problems and life applications to solve using the modulus equations and equalities.

### Key Terms

- ▶ Equation
- ▶ Inequality
- ▶ Graphical solution

### Materials

- ▶ Graphical calculator
- ▶ Graph paper
- ▶ Graphic programs

### First: Solving equations



#### Think and discuss


In one figure, represent the two curves of the two functions  $f$  and  $g$  where  $f$  is a modulus function and  $g$  is a constant function graphically.

Notice the graph, then answer:


- How many probable intersecting points are there for the two curves of the two functions together?
- Do the ordered pairs satisfy the rule of each function of both functions if the intersecting points of the two curves are found together?

#### Notice:


- At the intersecting points (if found),  $f(x) = g(x)$ , and vice versa for each  $x$  belong to the common domain of both functions.
- For any two functions  $f$  and  $g$ , the solution set of the equation  $f(x) = g(x)$  is the set of  $x$ -coordinates of the intersecting points of their two curves as shown in the following figures:



Solution set =  $\phi$



Solution set =  $\{a, b\}$



Solution set =  $\{a\}$

Solve the equation :  $|ax + b| = c$


### Example

- Solve the equation :  $|x - 3| = 4$  algebraically and graphically.

#### Solution

Let  $f(x) = |x - 3|$  and  $g(x) = 4$

- Graph the curve of the function  $f(x) = |x - 3|$  by translating the curve of  $f(x) = |x|$  3 units in the direction of  $\vec{ox}$



- 2) On the same figure, graph  $g(x) = 4$  where  $g$  is a constant function represented by a straight line parallel to  $x$ -axis and passing through point  $(0, 4)$   
 $\therefore$  The two points intersect at  $(-1, 4)$  and  $(7, 4)$   
 $\therefore$  The solution set is  $\{-1, 7\}$

**Algebraic solution :**

From the definition of the modulus function :  $f(x) = \begin{cases} x-3 & \text{when } x \geq 3 \\ -x+3 & \text{when } x < 3 \end{cases}$

When  $x > 3$  :  $x-3 = 4$  i.e. :  $x = 7 \in ]3, \infty[$

When  $x \leq 3$  :  $-x+3 = 4$  i.e. :  $x = -1 \in ]-\infty, 3]$

The solution set of the equation is  $\{-1, 7\}$  this matches with the graphical solution.

**Try to solve**

- 1) Solve each of the following equations graphically and algebraically.

a)  $|x| - 4 = 0$

b)  $|x| + 1 = 0$

c)  $|x-7| = 5$

**Some Properties of the Absolute Value**

**Learn**

- 1)  $|ab| = |a| \times |b|$  for example :

$|2 \times -3| = |-6| = 6$  and  $|2| \times |-3| = 2 \times 3 = 6$

- 2)  $|a + b| \leq |a| + |b|$

The equality happens only when **a and b** have the same sign. For example :

$|4 + 5| = |4| + |5| = 9$  et  $|-4 - 5| = |-4| + |-5| = 9$

- 3)  $|a - x| = |x - a|$

**Notice :**

- 1) If  $|x| = a$  then  $x = a$  or  $x = -a$  for each  $a \in \mathbb{R}^+$   
 2) If  $|a| = |b|$  then  $a = b$  or  $a = -b$  for each  $a \in \mathbb{R}, b \in \mathbb{R}$   
 3)  $|x|^2 = |x^2| = x^2$

**Solve the equation:  $la x + bl = lc x + dl$** 


**Example**

- 2) Solve the equation  $|x - 3| = |2x + 1|$  graphically.

**Solution**

Let  $f(x) = |x - 3|$  and  $g(x) = |2x + 1|$

The curve of  $f$  is the same curve of  $|x|$  by translation of a magnitude of 3 units in the direction of  $\vec{ox}$ .



$$\therefore g(x) = |2x + 1| = |2(x + \frac{1}{2})|$$

$$\therefore g(x) = 2|x + \frac{1}{2}|$$

The curve of  $g$  is the same curve of  $2|x|$  by horizontal translation of a magnitude of  $\frac{1}{2}$  unit in the direction of  $\overrightarrow{ox'}$  and the intersecting points of the two curves of the two functions  $f$  and  $g$  are  $(-4, 7)$  and  $(\frac{2}{3}, \frac{5}{2})$

The solution of the function is  $\{-4, \frac{2}{3}\}$

### Try to solve

2 Solve each of the following equations graphically.

a  $|x + 7| = |2x + 3|$

b  $|x - 2| + |x - 1| = \text{zero}$

### Example

3 Find the solution set for each of the following equation algebraically:

a  $|x + 7| = |x - 5|$

b  $\sqrt{x^2 - 6x + 9} = |9 - 2x|$

### Solution

a  $\therefore |x + 7| = |x - 5| \quad \therefore x + 7 = \pm(x - 5)$   
 $\therefore x + 7 = x - 5 \quad \text{but } 7 \neq -5 \text{ (impossible).}$   
 or  $x + 7 = -x + 5 \quad \text{i.e. } 2x = -2$   
 $\therefore x = -1 \quad \text{i.e. the solution set is } \{-1\}$

**Remember**  
 if  $a, b \in \mathbb{R}$  and  
 $|a| = |b|$   
 then  $a = \pm b$

### Check:

by substitution  $x = -1$  in the two sides of the equation, we find :  
 Right side = left side = 6. i.e. the solution set is  $\{-1\}$

b  $\therefore \sqrt{x^2 - 6x + 9} = |9 - 2x|$   
 $\therefore \sqrt{(x - 3)^2} = |9 - 2x| \quad \text{i.e. } \therefore |x - 3| = |9 - 2x|$   
 then  $x - 3 = \pm(9 - 2x)$   
 $\therefore x - 3 = 9 - 2x \quad , \quad 3x = 12 \quad \text{i.e. } x = 4$   
 or  $x - 3 = -9 + 2x \quad , \quad x = 6$

**Remember**  
 For any real number  
 a:  $\sqrt{a^2} = |a|$

By substituting the values of  $x$  in the two sides of the function

When  $x = 4$  right side = left side = 1  $\therefore x = 4$  is the solution of the equation

When  $x = 6$  right side = left side = 3  $\therefore x = 6$  is the solution of the equation

I.e. The solution set of the equation is :  $\{4, 6\}$

**P Try to solve**

3 Find the solution set for each of the following equation algebraically:

a  $|x - 1| - 2|2 - x| = 0$

b  $\sqrt{x^2 - 4x + 4} = 4$

**Second : Solving the inequalities**

You have previously learned the inequalities and known that they are mathematical phrases containing one of the symbols : ( $<$  ,  $>$  ,  $\leq$  ,  $\geq$ ). Solving the inequality means that you find the value or the set of values of the variable which satisfy the inequality and make it true.


**Solving the inequalities graphically**

The opposite figures shows a curve for each of the two functions  $f$  and  $g$  where:  $y_1 = f(x)$  and  $y_2 = g(x)$  and the solution set of the equation  $f(x) = g(x)$  is  $\{a, b\}$

**I.e.:**  $y_1 = y_2$  when  $x = a$  or  $x = b$

**We notice:**  $y_1 < y_2$  i.e.  $f(x) < g(x)$  when  $x \in ]a, b[$

$y_1 > y_2$  i.e.  $f(x) > g(x)$  when  $x \in ]-\infty, a[ \cup ]b, \infty[$


**Example**

**P Try to solve**

4 Find the solution set of the following inequalities using the graphs in example (7):

a  $|x + 2| \leq 2$

b  $|2x + 6| \leq 4$

c  $|x - 2| > 3$



## Solving the inequalities algebraically



## Learn

**First:** if  $|x| \leq a$  and  $a > 0$  then  $-a \leq x \leq a$

**Second:** if  $|x| \geq a$  and  $a > 0$  then  $x \geq a$  or  $x \leq -a$



## Example

5 Find the solution set of the following inequalities in the form of an interval:

a  $|x - 3| < 4$

b  $\sqrt{x^2 - 2x + 1} \geq 4$

## Solution

a  $\because |x - 3| < 4$  i.e.,  $-4 < x - 3 < 4$  Adding 3 to inequality

$\therefore -4 + 3 < x - 3 + 3 < 4 + 3$  i.e.,  $-1 < x < 7$

$\therefore$  the solution set =  $]-1, 7[$

b  $\because \sqrt{(x - 1)^2} = |x - 1|$  i.e.,  $|x - 1| \geq 4$

$\therefore x - 1 \geq 4$  ,  $x \geq 5$  or  $x - 1 \geq -4$  ,  $x \geq -3$

$x \in \mathbb{R} - ]-3, 5[$

$\therefore$  the solution set of the inequality is  $]-\infty, -3] \cup [5, \infty[$

## Remember

for  $a, b$  and  $c$   
 if:  $a < b, b < c$  then  
 $a < c$   
 if:  $a < b$  then  
 $a + c < b + c$   
 $a c < b c$  when  $c > 0$   
 $a c > b c$  when  $c < 0$

## Try to solve

5 Find the solution set of the following inequalities in the form of an interval :

a  $|x - 7| < 11$

b  $|3x + 7| \leq 8$

c  $\sqrt{x^2 - 6x + 9} \geq 8$

**Critical thinking:** Write down each of the following in the form of the absolute value inequality:

a  $4 \leq x \leq -4$

b  $x \leq -2, x \geq 2$

c  $0 < x < 6$

## Life applications



## Example Meteorology

6 A meteorological station has recorded the temperature of Cairo on a day. If the temperature has been  $32^\circ$  in difference  $7^\circ$  from its normal rate on that day. What is the expected temperature recorded in Cairo on that day?

## Solution

Let the temperature expected to be recorded in Cairo on that day =  $x^\circ$

$\therefore |x - 32| = 7$  i.e.  $x - 32 = \pm 7$

then  $x = 32 + 7 = 39$  or  $x = 32 - 7 = 25$   
 i.e. the temperature expected to be recorded is  $39^\circ$  or  $25^\circ$

**Try to solve**

- 6 **Athletic medicine** : Basim's weight differs from his ideal weight for 5kg. What is his probable weight if his ideal weight is 60 kg.?

**Example** **Want ads jobs**

- 7 A natural gas company employs the counter readers if their lengths range between 178 cm to 192 cm . Express the possible lengths to those who take up this job using the absolute value inequality.

**Solution**

Let the length of the person taking up the job =  $x$  cm


$$\therefore 178 \leq x \leq 192$$

By adding  $-185$  to the parts of the inequality :

$$178 - 185 \leq x - 185 \leq 192 - 185$$

$$-7 \leq x - 185 \leq 7$$

$$\text{i.e. } |x - 185| \leq 7$$



**Try to solve**

- 7 Write down the absolute value inequality which expresses the marks of a student in an exam that range between 60 to 100 marks :





**Activity**

**Using the functions in solving life and mathematical problems**

**Notice:** If a light ray falls on a reflective surface whose pathway is subjected to the modulus function . The measurement of incidence angle equals the measurement of reflection angle.

In addition , the pathway of the billiard ball before and after colliding it against the table edge.

The figure opposite illustrates that the billiard player kicks the black ball considering  $\vec{ox}$  and  $\vec{oy}$  the two perpendicular coordinates and the ball's pathway follows the curve of the function  $f$  where:  $f(x) = \frac{4}{3}|x - 5|$ . Does the black ball fall in pocket B? Explain mathematically.





## Exercises (1 - 5)



**Complete :**

- ① The solution set of the equation  $|x| = \frac{1}{2}$  is .....
- ② The solution set of the equation  $|x| + 3 = 0$  is .....
- ③ The solution set of the inequality  $|x - 2| \leq 0$  is .....

**Choose the proper solution set from the following list for each equation or inequality :**

- ④  $|x - 2| = 3$
- ⑤  $|x - 2| < 3$
- ⑥  $|x - 2| > -3$
- ⑦  $|x - 2| \leq 3$
- ⑧  $|x - 2| > 3$
- ⑨  $|x - 2| = -3$

- |   |                        |
|---|------------------------|
| a | $]-1, 5[$              |
| b | $\mathbb{R}$           |
| c | $\{-1, 5\}$            |
| d | $\mathbb{R} - [-1, 5]$ |
| e | $\emptyset$            |
| f | $[-1, 5]$              |

**Find the solution set for each of the following equations algebraically :**

- |                       |                        |                             |
|-----------------------|------------------------|-----------------------------|
| ⑩ $ x + 3  = 6$       | ⑪ $ 2x - 7  = 5$       | ⑫ $ 3 - 2x  = 7$            |
| ⑬ $ x - 3  =  x + 1 $ | ⑭ $ 2x + 1  =  x - 3 $ | ⑮ $\sqrt{x^2 - 2x + 1} = 4$ |

**Find the solution set for each of the following equations graphically :**

- |                 |                       |                  |
|-----------------|-----------------------|------------------|
| ⑯ $ x + 4  = 3$ | ⑰ $ x - 1  =  x + 3 $ | ⑱ $ 2x - 5  = 3$ |
|-----------------|-----------------------|------------------|

**Find the solution set for each of the following equations graphically:**

- |                 |                    |                 |
|-----------------|--------------------|-----------------|
| ⑲ $ x - 1  < 3$ | ⑳ $ x - 2  \leq 5$ | ㉑ $ x + 3  > 2$ |
|-----------------|--------------------|-----------------|

**Find the solution set for each of the following equations algebraically:**

- |                  |                     |                     |
|------------------|---------------------|---------------------|
| ㉒ $ 2x - 1  > 3$ | ㉓ $ 2x + 3  \leq 7$ | ㉔ $ 3x - 7  \geq 2$ |
|------------------|---------------------|---------------------|

⑲ **Roads :** two roads; the first road is represented by the curve of the function  $f$  where  $f(x) = |x - 4|$  and the second one is represented by the curve of the function  $g$  where  $g(x) = 3$ . If the two roads get intersected at the two points A and B, find the distance from A to B known that the length unit represents 1 km only.

⑳ Write down the absolute value inequality which expresses the temperature measured by the medical thermometer and ranges between  $35^\circ$  and  $42^\circ$ .

## Unit summary

- 1 The function :** is a relation between two non-null sets  $X$  and  $Y$  so that each element in  $x$  has one and only one element of  $Y$  and the function is symbolically written in the form  $f: X \longrightarrow Y$ . The function is determined by the three elements; the domain, co-domain and the rule of the function.

The function  $f$  is called a real function if each of its domain and co-domain are the set of the real numbers or a subset of it.

- 2 The vertical line test :** If a relation is represented by a set of points in a orthogonal coordinate plane and the vertical line intersects its graphical representation at each element of the domain elements at one point only, then the relation represents a function.

- 3 Piecewise- defined function :** is a real function in which each subset of its domain has a different definition rule.

- 4 Monotony of function :** the function  $f$  is **increasing** in the interval  $]a, b[$  if each of  $x_1$  and  $x_2 \in ]a, b[$  and  $x_2 > x_1$ , then  $f(x_2) > f(x_1)$ :

The function  $f$  is **decreasing** in the interval  $]a, b[$  if each of  $x_1, x_2 \in ]a, b[$  when  $x_2 > x_1$ , then  $f(x_2) < f(x_1)$

and the function is **constant** in the interval  $]a, b[$  if each  $x_1, x_2 \in ]a, b[$  and  $x_2 > x_1$ , then  $f(x_2) = f(x_1)$

- 5 Even and odd functions :**

**Even function :** it is said the function  $f: x \longrightarrow y$  is an even function if  $f(-x) = f(x)$  for each  $x$  and  $-x \in X$ .

**Odd function :** it is said the function  $f: x \longrightarrow y$  is an odd function if  $f(-x) = -f(x)$  for each  $x$  and  $-x \in X$ .

### Important properties :

If each of  $f_1$  and  $f_2$  are even functions and each of  $g_1$  and  $g_2$  are odd functions, then :

- |  |  |
|--|--|
| <b>1)</b> $f_1 + f_2$ even function      | <b>2)</b> $g_1 + g_2$ odd function.                  |
| <b>3)</b> $f_1 \times f_2$ even function | <b>4)</b> $g_1 \times g_2$ even function.            |
| <b>5)</b> $f_1 \times g_2$ odd function  | <b>6)</b> $f_1 + g_2$ neither even nor odd function. |

- 6 Linear function :** its simplest form is  $f(x) = x$ , represented by a straight line passing through the point  $(0, 0)$  and its slope = 1

- 7 Quadratic function :** its simplest form is  $f(x) = x^2$ , the curve vertex point is  $(0, 0)$  and the equation of the symmetry axis  $x = 0$ .

- 8 Cubic function :** its simplest form is  $f(x) = x^3$  and the symmetry point of its curve is  $(0, 0)$ .

- 9 Modulus function (absolute value function)**

The simplest form of the modulus function is  $f(x) = |x|$  and defined as follows:

$$f(x) = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

It is represented by two rays starting from the point of  $(0, 0)$ . The slope of the first ray = 1

and the other = -1, therefore  $|x| \geq 0$ ,  $|-x| = |x|$  and  $\sqrt{x^2} = |x|$

- 10 Rational function :** its simplest form is  $f(x) = \frac{1}{x}$  and the symmetry point of its curve is  $(0, 0)$ .
- 11 Geometrical transformations** of the function  $f$  where  $y = f(x)$  and  $a > 0$  are determined as follows :
- If  $y = f(x) + a$  it is represented by translating the curve of  $f$  in the positive direction of y-axis in a magnitude of  $a$
  - If  $y = f(x) - a$  it is represented by translating the curve of  $f$  in the negative direction of y-axis in a magnitude of  $a$
  - If  $y = f(x + a)$  it is represented by translating the curve of  $f$  in the negative direction of x-axis in a magnitude of  $a$
  - If  $y = f(x - a)$  it is represented by translating the curve of  $f$  in the positive direction of x-axis in a magnitude of  $a$ .
  - If  $y = f(x)$  it is represented by reflection of the curve of  $f$  in x-axis.
  - If  $y = af(x)$  it is represented by the stretching the two vertices of the curve  $f$  if  $a > 1$  and by shrinking the two vertices of the curve if  $0 < a < 1$ .
- 12 Properties of absolute value :**
- a**  $|a \cdot b| = |a| \times |b|$
  - b**  $|a + b| \leq |a| + |b|$
  - c** If  $|x| \leq a$ ,  $a > 0$  then  $-a \leq x \leq a$
  - d** If  $|x| \geq a$ ,  $a > 0$  then  $x \geq a$  or  $x \leq -a$
- 13) Solving the equation :** for any two functions  $f$  and  $g$ , the solution set of the function  $f(x) = g(x)$  is the set of the x coordinates of the intersecting points of their two curves.
- 14) Solving the inequality:** is to find the set of the values of the variable which make the inequality true.



### General Exercises

For more exercises please visit the website of the Ministry of Education.



### Enrichment Information

Please visit the following link.





## Accumulative test



- 1 Graph the two curves of the functions  $f$  and  $g$  where  $f(x) = x + 1$  and  $g(x) = 5 - x$ , then find:
- The two coordinates of the intersecting points of each curve with  $x$ -axis.
  - The two coordinates of the intersecting point of the two curves.
  - Area of the triangle determined by the two intersected straight lines and  $x$ -axis.
- 2 Use the curve of the function  $f$  where  $f(x) = |x|$  to represent the function  $g$  where  $g(x) = |x - 1| - 2$ , then find the range of the function  $g$ .
- 3 Graph the curve of the function  $f$  where  $f(x) = \begin{cases} x^2 & \text{for each } -2 \leq x < 2 \\ 6 & \text{for each } 2 \leq x \end{cases}$
- Then determine the range of the function and investigate its monotony.

- 4 In the figure opposite :

- Write the coordinate of the curve vertex point.
- Write the rule of the function.
- Find the range of the function and investigate its monotony.
- Write the equation of the symmetry axis.



- 5 Graph the curve of the function  $f$  where  $f(x) = (x - 1)^3$  and deduce the range and monotony of the function, then tell whether it is even, odd or otherwise.
- 6 Use the curve of the function  $f$  where  $f(x) = \frac{1}{x}$  to represent the function  $g$  where  $g(x) = f(x) + 2$  then write the symmetry point of the resulted function and investigate its monotony.
- 7 If the function  $f$  where  $f(x) = \frac{1}{x+1}$ , find the domain of the function  $f$  and the symmetry point of the curve of this function.

Solve the equation  $f\left(\frac{1}{x}\right) = 4$

- 8 Find the solution set for each of the following graphically

**a**  $|x - 4| = 3$                       **b**  $|x - 4| \geq 3$

- 9 Find the solution set of the following equations and inequalities algebraically :

**a**  $|x + 5| = 9$                       **b**  $\sqrt{4x^2 - 12x + 9} = |x + 1|$                       **c**  $|2x - 5| \leq 7$

**d**  $|3x + 1| > 7$



## Unit Two

# Exponents, Logarithms and their Applications



### Unit introduction

The concept of logarithms had been used in mathematics early the Seventeenth century by the scientist John Napier as a method to simplify calculations to help navigators, scientists, engineers and others depend on to do their calculations easier by using the calculating ruler and logarithmic tables. They had benefited from the properties of logarithms by substituting the multiplying operations to find the logarithm of the product of two numbers, by using the property of addition with respect to the property of  $\log_a(xy) = \log_a x + \log_a y$ . Thanks to the scientist Leonhart Euler. In the eighteenth century, he connected the concept of logarithm with the concept of the exponential function to widen the concept of logarithms and connect with the functions.

Logarithmic scaler can be used in various fields. For example, the decibel is a logarithmic unit used to measure the sound intensity and voltage ratio. Logarithmic scaler is also used to measure the power of hydrogen Ph( it is a logarithmic scaler) to identify the acidity of a solution in chemistry.



### Unit objectives

**By the end of this unit, the student should be able to:**

- Identify the exponential function.
- Identify the graphical representation of the exponential function and deduce its properties.
- Identify the rules of the rational exponents.
- Solve an exponential equation in the form:  $a^x = b$ .
- Identify the logarithmic equation.
- convert from the exponential form into logarithmic form algebraically and vice versa.
- I identify the graphical representation of the logarithmic function in limited intervals and deduce its properties.
- Deduce the relation between the exponential and logarithmic functions graphically.
- Identify the rules of logarithms.
- Solve logarithmic equations.
- Solve problems including applying the rules of logarithms.
- Identify the common logarithms of base 10.
- Find the value of logarithms using the calculator.
- Use the calculator to solve some exponential equations.



## Key - terms

- The  $n^{\text{th}}$  Power
- Base
- Exponent
- $n^{\text{th}}$  Root
- Rational - exponent

- Exponential Function
- Exponential Growth
- Exponential Decay
- Domain
- Range

- Reflection
- Logarithm
- Logarithmic Equation
- Logarithmic Function


## Unit lessons

- Lesson 1:** Rational exponents
- Lesson 2:** Exponential Function and its applications
- Lesson 3:** Solving exponential equations
- Lesson 4:** Logarithmic function and its graphical representation
- Lesson 5:** Some properties of logarithms

## Materials

Scientific calculator - geogebra-graph

## Unit planning guide



**You will learn**

- Generalize the rules of exponents.
- $n^{\text{th}}$  root.
- Rules of rational exponents.

**Key terms**

- The  $n^{\text{th}}$  power
- Base
- Exponent
- $n^{\text{th}}$  root
- Rational exponent

**Materials**

- Scientific calculator
- Graphic programs

**Preface**

You have previously learned the square roots of a non-negative real number and identified some properties of both cubic and square roots. You have also learned the integers and identified some of their own properties. In this lesson, you are going to learn the rational exponents.

**Learn****Integer exponents:**

**Definition**

- 1) For each  $a \in \mathbb{R}$  and for each  $n \in \mathbb{Z}$ , then:

$a^n = a \times a \times a \times \dots \times a$  (where factor  $a$  is repeated  $n$  times).

$(a^n)$  is called the  $n^{\text{th}}$  power of  $a$ , where  $a$  is called the base and  $n$  is called the exponent. We say  $a$  exponent  $n$ .

1

- 2)  $a^0 = 1$  for all  $a \in \mathbb{R} - \{0\}$

- 3)  $a^{-1} = \frac{1}{a}$ ,  $a^{-n} = \frac{1}{a^n}$   $a \neq 0$

**Properties of Integer Exponents:**

For each  $m$  and  $n \in \mathbb{Z}$ ,  $a, b \in \mathbb{R}$ ,  $b \neq 0$ , then:

$$\triangleright a^m \times a^n = a^{m+n} \qquad \triangleright (ab)^n = a^n b^n$$

$$\triangleright \frac{a^m}{a^n} = a^{m-n} \qquad \triangleright \left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$$

$$\triangleright (a^m)^n = a^{m \cdot n}$$

**Example**

- 1 Find the following term in the simplest form:  $\frac{(8)^{-3} \times (18)^2}{81 \times (16)^{-3}}$

**Solution**

$$\begin{aligned} \text{The expression} &= \frac{(2^3)^{-3} \times (2 \times 3^2)^2}{3^4 \times (2^4)^{-2}} = \frac{2^{-9} \times 2^2 \times 3^4}{3^4 \times 2^{-8}} \\ &= 2^{-9+2+8} \times 3^{4-4} \\ &= 2^1 \times 3^0 = 2 \times 1 = 2 \end{aligned}$$

**P Try to solve**

① Find the value of the following term in the simplest form:  $\frac{(27)^3 \times (12)^2}{16 \times (81)^2}$

② Prove that:  $\frac{2^6 \times 9^{4-1}}{3 \times 18^4} = 3$



**Learn**

**The  $n^{\text{th}}$  Root**

You have known that the square root of a number is an inverse operation of squaring this number. Similarly, the  $n^{\text{th}}$  root of a number is an inverse operation to place the  $n^{\text{th}}$  power of this number.

**Example:**

1 If  $x^3 = 8$  then 2 is the cubic root of 8 i.e.  $\sqrt[3]{8} = 2$

2 If  $x^5 = 32$  then 2 is the fifth root of 32 i.e.  $\sqrt[5]{32} = 2$

3 If  $x^n = a$  then  $x$  is the  $n^{\text{th}}$  root of  $a$  i.e.  $\sqrt[n]{a} = x$

Definition

For any real number,  $a \geq 0$ ,  $n \in \mathbb{Z}^+ - \{1\}$ , then  $a^{\frac{1}{n}} = \sqrt[n]{a}$   
This relation is true when  $a < 0$  and  $n$  is an odd number  $> 1$

**Example:**

1

$(16)^{\frac{1}{4}} = \sqrt[4]{16} = 2$        $(-9)^{\frac{1}{2}} = \sqrt{-9} \notin \mathbb{R}$

$-(27)^{\frac{1}{3}} = -\sqrt[3]{27} = -3$        $(-243)^{\frac{1}{5}} = \sqrt[5]{-243} = -3$



**Notice**

Index of the root  $\sqrt[n]{a}$  symbol of the root  
number inside the root



**Example**

② If  $x^n = a$ , find the values of  $x$  in  $\mathbb{R}$  (if found) in each of the following cases:

a  $n = 5$ ,  $a = \text{zero}$

b  $n = 4$ ,  $a = 81$

c  $n = 2$ ,  $a = -4$

d  $n = 3$ ,  $a = -8$

**Solution**

a when  $n = 5$ ,  $a = \text{zero}$  then  $x^5 = 0$  then  $x = \sqrt[5]{0} = 0$

b when  $n = 4$ ,  $a = 81$  then  $x^4 = 81$  then  $x = \pm \sqrt[4]{81} = \pm 3$

c when  $n = 2$ ,  $a = -4$  then  $x^2 = -4$  then  $x = \pm \sqrt{-4} \notin \mathbb{R}$

d when  $n = 3$ ,  $a = -8$  then  $x^3 = -8$  then  $x = \sqrt[3]{-8} = -2$

from the previous example, we deduce that:

if  $x^n = a$ , then the values of  $x$ , which satisfy the equation become clear in the following table:

$n$	$a$	$\sqrt[n]{a}$
$n \in \mathbb{Z}^+ - \{1\}$	$a = 0$	$\sqrt[n]{a} = \text{zero}$
positive even integer	$a > 0$	there are two real roots $\pm \sqrt[n]{a}$
positive even integer	$a < 0$	there are no real roots.
positive odd integer, $n \neq 1$	$a \in \mathbb{R}$	there is only one real root $\sqrt[n]{a}$

**P Try to solve**

3 Find the values of  $x$  in each of the following (if possible):

a  $x^2 = 36$

b  $x^5 = -32$

c  $x^3 = 125$

d  $x^4 = 1296$

e  $x^2 = -49$

f  $x^7 = -128$

4 **Critical thinking:** use a numerical example to show the difference between the sixth root of  $a$  and  $\sqrt[6]{a}$ .

Definition

if  $n \in \mathbb{Z}^+ - \{1\}$ ,  $m \in \mathbb{Z}^+$ ,  $\sqrt[n]{a} \in \mathbb{R}$  then:  $a^{\frac{m}{n}} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m$

**Example:**

$$(16)^{\frac{3}{2}} = (\sqrt{16})^3 = (4)^3 = 64$$

$$\sqrt[3]{(-125)^2} = (\sqrt[3]{-125})^2 = (-5)^2 = 25$$

**Example**

3 Find each of the following in the simplest form:

a  $-\sqrt[3]{8a^6b^9}$

b  $\pm\sqrt{64(a^2+3)^6}$

**Solution**

a  $-\sqrt[3]{8a^6b^9} = -\sqrt[3]{(2a^2b^3)^3} = -2a^2b^3$

b  $\pm\sqrt{64(a^2+3)^6} = \pm\sqrt{[8(a^2+3)^3]^2}$   
 $= \pm 8(a^2+3)^3$

**Try to solve**

5 Find each of the following in the simplest form:

a  $\sqrt[4]{625a^{12}}$

b  $\sqrt[3]{-243b^5}$

c  $\sqrt[3]{128(a+b)^7}$

### Using The Modulus

the modulus of a number is used if the index of the root ( $n$ ) is an even number, then  $\sqrt[n]{a^2} = |a|$ , but if the index is an odd number, it is not necessary to use the modulus.

$$\sqrt[n]{x^2} = \begin{cases} |x| & \text{if } n \text{ is even.} \\ x & \text{if } n \text{ is odd.} \end{cases}$$



#### Example

4 Find each of the following in the simplest form :

a  $\sqrt{9x^2}$

b  $\sqrt[3]{-8x^3}$

c  $\sqrt[4]{(2-\sqrt{3})^4}$

d  $\sqrt[4]{(1-\sqrt{7})^4}$

#### Solution

a  $\sqrt{9x^2} = \sqrt{(3x)^2} = |3x|$

b  $\sqrt[3]{-8x^3} = \sqrt[3]{(-2x)^3} = -2x$

c  $\sqrt[4]{(2-\sqrt{3})^4} = |2-\sqrt{3}| = 2-\sqrt{3}$  where  $2 > \sqrt{3}$

d  $\sqrt[4]{(1-\sqrt{7})^4} = |1-\sqrt{7}| = \sqrt{7}-1$  where  $\sqrt{7} > 1$



#### Notice

the square of any of the two numbers (a) or (-a) is  $a^2$

#### Try to solve

6 Find each of the following in the simplest form:

a  $\sqrt[4]{16a^{12}}$

b  $\sqrt[3]{(x-2)^{18}}$

c  $\sqrt[4]{(2-\sqrt{5})^4}$

d  $\sqrt[4]{(2-\sqrt{5})^4}$

#### Definition

if  $n \in \mathbb{Z}^+ - \{1\}$ ,  $m \in \mathbb{Z}^+$ ,  $\sqrt[n]{a} \in \mathbb{R}$  then:  $a^{\frac{m}{n}} = \frac{1}{a^{\frac{1}{n}}}$

#### 4

Example:  $7^{\frac{3}{5}} = \frac{1}{7^{\frac{2}{5}}}$ ,  $\frac{1}{4^{\frac{2}{3}}} = 4^{\frac{1}{3}}$


#### Definition

if  $n \in \mathbb{Z}^+ - \{1\}$ ,  $\sqrt[n]{a}$ ,  $\sqrt[n]{b}$  are two real numbers, then:

>  $\sqrt[n]{ab} = \sqrt[n]{a} \times \sqrt[n]{b}$

>  $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$  where  $b \neq \text{zero}$

#### 5

 **Example**

5 Find each of the following in the simplest form:

a  $\frac{\sqrt{8} \times 4^{-1} \times 2^{\frac{3}{2}}}{6^{-2} \times 3^2}$

b  $\frac{32^{-\frac{3}{5}} \times 8^{\frac{2}{5}}}{\sqrt[3]{4} \times \sqrt[3]{16^5}}$

 **Solution**

a the expression =  $\frac{8^{\frac{1}{2}} \times 4^{-1} \times 2^{\frac{3}{2}}}{6^{-2} \times 3^2}$   
 $= \frac{(2^3)^{\frac{1}{2}} \times (2^2)^{-1} \times 2^{\frac{3}{2}}}{(3 \times 2)^{-2} \times 3^2}$   
 $= \frac{(2^{\frac{3}{2}} \times 2^{-2} \times 2^{\frac{3}{2}})}{3^{-2} \times 2^{-2} \times 3^2}$   
 $= 2^{\frac{3}{2} - 2 - \frac{3}{2}} \times 2 \times 3^{2 - 2}$   
 $= 2^{\text{zero}} \times 3^{\text{zero}} = 1$

converting roots into rational exponents.

factorizing each base into its primary factors.

By simplifying

b the expression =  $\frac{32^{-\frac{3}{5}} \times 8^{\frac{2}{5}}}{4^{\frac{1}{3}} 16^{\frac{5}{3}}}$   
 $= \frac{(2^5)^{-\frac{3}{5}} (2^3)^{\frac{2}{5}}}{(2^2)^{\frac{1}{3}} (2^4)^{\frac{5}{3}}}$   
 $= \frac{2^{-3} \times 2^2}{2^{\frac{1}{3}} \times 2^{\frac{20}{3}}}$   
 $= 2^{3 \times 2 - \frac{1}{3} - \frac{20}{3}} = 2^2 = 4$

converting roots into rational exponents.

factorizing each base into its primary factors.

 **Try to solve**

7 Find each of the following in the simplest form :

a  $\frac{\sqrt[3]{243} \times \sqrt[3]{8^{-1}}}{\sqrt{2} \times \sqrt[3]{9}}$

b  $\frac{\sqrt[3]{4} \times \sqrt[3]{2}}{\sqrt{2} \times \sqrt[3]{4}}$

**Solving the equations:**
 **Example**

6 Find the solution set for each of the following equations in R:

a  $x^{\frac{2}{3}} = 9$

b  $(x + 1)^{\frac{3}{4}} = 8$

 **Solution**

a  $\therefore x^{\frac{2}{3}} = 9$  By placing the third power of the two sides

$\therefore (x^{\frac{2}{3}})^3 = 9^3$

$\therefore x^2 = 9^3$  By taking off the square root of both sides

$\sqrt{x^2} = \sqrt{9^3} \quad \therefore |x| = 27 \quad \therefore x = \pm 27$

 $\therefore$  The solution set =  $\{27, -27\}$

- b**  $(x + 1)^{\frac{1}{4}} = 8$  By placing the fourth power of the two sides

$$\therefore (x + 1)^{\frac{1}{4} \times 4} = 8^4$$

$$\therefore (x + 1) = (\sqrt[4]{8^4})^4$$

$$\therefore x + 1 = 2^4$$

$$\therefore x = 15$$

$$\therefore \text{The solution set} = \{15\}$$

**P Try to solve**

- 8** Find the solution set for each of the following equations in R:

**a**  $x^{\frac{5}{2}} = 32$

**b**  $\sqrt[5]{(x-1)^5} = \frac{1}{32}$

**Example**

- 7 Geometry:** If the side length of the square whose area  $m$  is given by the relation  $L = m^{\frac{1}{2}}$

- a** Calculate the side length of the square whose area is  $25\text{cm}^2$ .  
**b** Calculate the side length of the square whose area is  $17\text{cm}^2$  approximating the result to one decimal.

**Solution**

**a**  $L = 25^{\frac{1}{2}} = \sqrt{25} = 5 \text{ cm}$


**b**  $L = 17^{\frac{1}{2}} = \sqrt{17} \approx 4.12310$

By approximating to one decimal

$\therefore L \approx 4.1 \text{ cm}$

**P Try to solve**

- 9** If  $L$  is the side length of a cub whose volume  $P$  is given by the relation  $L = P^{\frac{1}{3}}$ , find the side length of the cube whose volume is 27.

 **Exercises 2 - 1** 

- 1** Write down each of the following in an exponential form:

**a**  $\sqrt{x}$

**b**  $\sqrt[3]{a^3}$

**c**  $2\sqrt{n}$

**d**  $\sqrt[4]{a^2 b^3}$

**e**  $\sqrt[5]{x^5}$

**f**  $\frac{\sqrt{x}}{\sqrt[3]{x^2}}$

- 2** Write down each of the following in a root form:

**a**  $a^{\frac{1}{2}}$

**b**  $b^{\frac{2}{3}}$

**c**  $6y^{\frac{3}{4}}$

**d**  $8b^{\frac{4}{9}}$

**e**  $(3x)^{-\frac{2}{3}}$

**f**  $5^{\frac{1}{2}}$

- 3** Find the value of each of the following in the simplest form:

**a**  $(16)^{\frac{3}{4}}$

**b**  $(-32)^{\frac{2}{3}}$

**c**  $27^{-\frac{4}{3}}$

**d**  $(\frac{1}{8})^{\frac{2}{3}} + (\frac{1}{4})^{\frac{1}{2}}$

**e**  $\frac{\sqrt[3]{4}}{\sqrt{2}}$

**f**  $\frac{1}{(2^{-2} \times 4^{\frac{1}{2}} \times 8^{\frac{2}{3}})^{-2}}$

4 Find in the simplest form:

a  $(a^{-\frac{2}{3}})^{-3}$

b  $\sqrt[4]{x} \times x^{\frac{1}{2}}$

c  $(3^2 + 4^2)^{\frac{1}{2}}$

d  $(x^{\frac{1}{2}} + y^{\frac{1}{2}})(x^{\frac{1}{2}} - y^{\frac{1}{2}})$

e  $(x^{\frac{1}{3}} - y^{\frac{1}{3}})(x^{\frac{2}{3}} + x^{\frac{1}{3}}y^{\frac{1}{3}} + y^{\frac{2}{3}})$

f  $(x^{\frac{1}{2}} + x^{-\frac{1}{2}})^2$

g  $(1^3 + 2^3 + 3^3)^{\frac{1}{2}}$

5 Reduce each of the following in the simplest form:

a  $\sqrt[3]{243} + \sqrt[3]{512}$

b  $(\frac{16}{81})^{\frac{1}{3}} \times (\frac{729}{8})^{\frac{1}{2}}$

c  $(16)^{\frac{3}{2}} \div (8)^{\frac{3}{2}}$

d  $(27)^{\frac{2}{3}} - (64)^{\frac{5}{6}}$

e  $\sqrt{0.1} \times \sqrt[3]{0.216} \times \sqrt{2.5}$

f  $\frac{8^{\frac{3}{8}} \times 4^{\frac{3}{16}}}{2^{-\frac{3}{4}}}$

g  $(125)^{\frac{2}{3}} \times 81^{\frac{1}{4}} \times (15)^{-1}$

h  $\frac{16^{x-1} \times 9^{x+1}}{8^{x-1} \times 18^{x+2}}$

Choose the correct answer:

 6 If  $\sqrt{x^2} = 9$ , then  $x \in$ 

a  $\{27\}$

b  $\{27, -27\}$

c  $\{1\}$

d  $\phi$

 7  $64^{-\frac{1}{6}} =$ 

a 2

b -2

c  $\frac{1}{2}$

d  $-\frac{1}{2}$

 8  $\sqrt[3]{x^{-3}} =$ 

a  $x^{-1}$

b  $-x$

c  $|x^{-1}|$

d  $-|x|$

 9  $\sqrt[3]{x^4 y^6} =$ 

a  $x y^2$

b  $\pm x y^2$

c  $|x|y^2$

d  $|x|y^2|$

 10 If  $x^{-\frac{3}{2}} = 8$ , then  $x =$ 

a 4

b -4

c  $\frac{1}{4}$

d  $-\frac{1}{4}$

 11  $\frac{6^{-\frac{1}{3}} \times 6^{\frac{3}{3}}}{\sqrt[3]{36}} =$ 

a 1

b 6

c  $\frac{1}{6}$

d  $\sqrt{6}$



- 12 Find the solution set for each of the following equations in R:

a  $x^{\frac{1}{2}} = 5$

b  $x^{\frac{7}{2}} = \frac{1}{128}$

c  $\sqrt{x^3} = 27$

d  $(x - 5)^{\frac{5}{2}} = 32$

e  $3x^{-\frac{3}{4}} = \frac{3}{8}$

f  $2^{3x-1} = \frac{16}{\sqrt{2}}$

- 13 **Economy:** if it is known that the profit ( $r$ ) of a bank on a sum of money ( $a$ ) after ( $n$ ) year is given by the relation  $r = \left(\frac{c}{a}\right)^{\frac{1}{n}} - 1$  where  $c$  is the total money after  $n$  year. If Gamal has deposited 10000 LE and after 3 years the sum of money becomes 12597, find the yearly percentage of the profit.

- 14 **Discover the error:**

a if  $x^{\frac{3}{2}} = 4$ , then  $x = 8$

b  $\sqrt[4]{x^4} = x$

- 15 **Reduce the term:**  $\frac{\sqrt{a}}{a\sqrt{a}}$

- 16 **Activity:**

use the calculator to simplify the following operations (Round to tow decimals):

a  $75(1.21)^{\frac{19}{2}}$

b  $\frac{\sqrt[3]{2^{-1}} \times \sqrt[4]{7^{-2}}}{\sqrt{4^{-3}}}$

- 17 **Trade:** Mohammed has started a project to grow rabbits, if the number of rabbits at the beginning of the project was 75 rabbits and the number of rabbits in their reproduction has followed the relation  $Z = 75(4.22)^{\frac{n}{6}}$  where  $n$  is the number of months. Find the number of rabbits expected over 5 months..

- 18 **Volumes:** If the side length of a cube  $L$  is determined by the relation  $L = \sqrt[3]{V}$  where  $V$  is the volume of the cube in cubic units. Find the side length of a cube whose volume is  $1331\text{cm}^3$ .

### Creative thinking

#### Volumes :

- 19 if the radius length of a sphere is  $r$  whose volume  $V$  is given by the relation  $r = \sqrt[3]{\frac{3V}{4\pi}}$  :

A Find the radius length of a sphere whose volume is  $27000\text{cm}^3$ .

B calculate the change in the volume of the sphere as its radius length increases twice.

**You will learn**

- Exponential function.
- Representing the exponential function graphically.
- Properties of the exponential function.

**Key terms**

- Exponential Function
- Exponential Growth
- Exponential Decay

**Materials**

- Scientific calculator
- Graphic programs

**Add to your information**

the exponential function  $f(x) = a^x$  in case  $a > 1$  is called the growth function and is related to many life applications such as over population, and the banking compound interest. The exponential function  $f(x) = a^x$  in case  $0 < a < 1$  is called the decay function and is related to many applications such as the half-life of the radioactive atoms.

**Preface**

In our daily life, there are a lot of situations that need very accurate calculations such as banking profits, over population, cell reproduction in some organisms and the half-life of the radioactive atoms and so on. These situations require the concept of the exponential function which we are going to learn and investigate some of its properties.

**Learn****Exponential Function****Definition**

if  $a$  is a positive real number  $a \neq 1$  then the function:

$$f \text{ where } f: \mathbb{R} \rightarrow \mathbb{R}^+, f(x) = a^x$$

is called an **exponential function** whose base is  $a$

**Notice**

**In algebraic function:** the independent variable ( $x$ ) is the base while the power is a real number.

**In exponential function:** The independent variable ( $x$ ) is the power while the base is the real number and does not equal one.

**Verbal expression:** Explain why the function  $f(x) = (-3)^x$  where  $x \in \mathbb{R}$  is an exponential function.

**Graphical Representation of the Exponential Function****Example**

- use the values of  $x \in [-3, 3]$  to graph in one figure a part of the curve of each of the following two functions:


$$f(x) = 2^x, \quad g(x) = \left(\frac{1}{2}\right)^x$$

**Solution**

$x$	-3	-2	-1	0	1	2	3
$f(x)$	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	4	8
$g(x)$	8	4	2	1	$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{8}$

we can deduce the next properties of the exponential function from the graph:

- the function  $f: f(x) = 2^x$  is increasing on its domain because ( $a > 1$ )  
the function  $g: g(x) = \left(\frac{1}{2}\right)^x$  is decreasing on its domain because ( $0 < a < 1$ )
- the range of both functions is  $\mathbb{R}^+$



- 3 the curve of the function  $f: f(x) = 2^x$  is the image of the function curve  $g: g(x) = (\frac{1}{2})^x$  by reflection on y-axis

**Try to solve**

- 1 Use the values of  $x \in [-2, 2]$  to graph in one figure the curve of each of the functions  $f_1(x) = 2^x$ ,  $f_2(x) = 3^x$  and  $f_3(x) = 4^x$ .

**Example**

- 2 if  $f(x) = 3^x$ , complete the following :

a  $f(2) = \underline{\hspace{2cm}}$       b  $f(x+2) = \underline{\hspace{2cm}} \times 3^x$       c  $f(x) \times f(-x) = \underline{\hspace{2cm}}$

**Solution**

a  $f(2) = 3^2 = 9$       b  $f(x+2) = 3^{x+2} = 3^x \times 3^2 = 9 \times 3^x$   
 c  $f(x) \times f(-x) = 3^x \times 3^{-x} = 3^{x-x} = 3^{\text{zero}} = 1$

**Application on the exponential function:**

**First : The exponential growth**

We can use the function  $f$  where  $f(n) = a(1+g)^n$  to represent the exponential growth of a quantity  $a$  by a constant percentage  $g$  in equal periods which their number is  $n$ . ( discuss with your teacher how to deduce this relation).

**The compound interest**

To calculate the total sum  $C$  of an amount of money  $a$  invested in a bank giving an annually compound interest  $r$  ( percentage) for  $n$  number of year in division periods of the annual return into  $x$  period , then the sum of money is given by the relation :

$$C = a \left(1 + \frac{r}{x}\right)^{nx}$$

- 3 A man has deposited 5000LE in a bank giving an annually compound interest 8%. Find the sum of the money after 10 years in each of the following cases:  
 A annual interest ,      B three months' interest.      C Monthly interest.

**Solution**

By using the relation  $C = a \left(1 + \frac{r}{x}\right)^{nx}$  where  $x$  is the annual division :

a annual interest       $\therefore x = 1$   
 $c = 5000 (1 + 0.08)^{10} = 10794.62$  L.E  
 b three months' interest       $\therefore x = 4$   
 $c = 5000 \left(1 + \frac{0.08}{4}\right)^{10 \times 4} = 11040.2$  L.E

- c Monthly interest  $\therefore x = 12$


$$c = 5000 \left(1 + \frac{0.08}{12}\right)^{10 \cdot 12} = 11098.2 \text{ L.E}$$

**P Try to solve**

- 2 A man has deposited 1000LE in a bank giving a yearly complex profit 5%. Find the sum of the money after 8 years in each of the following cases:
- A yearly interest .      B 6 months' interest.      C Monthly interest.

### Second : Exponential Decay

the function  $f: f(n) = a(1 - r)^n$  whose base is lesser than one and greater than zero can be used to represent the exponential decay in a constant percentage  $g$  in equal periods whose number is  $n$ .

 **Example**

- 4 If the maximal production of a gold mine is 1850 kg per year and this production starts to decrease yearly in ratio 9%:
- A write an exponential function representing the gold production of this mine after  $n$  year.
- B Estimate the production of this mine after 8 years to the nearest kg.

 **Solution**

$$a = 1850 \quad , \quad r = 0.09$$

a the function of the exponential decay  $f(n) = (a - r)^n$   
 $f(n) = 1850 (1 - 0.09)^n$

b After 8 years (by substituting  $n = 8$ )  
 $\therefore f(8) = 1850 (1 - 0.09)^8 \simeq 870 \text{ kg}$

**P Try to solve**

- 3 if the marketing price of a car decreases according to the relation  $x = 150000(0.94)^n$ , where  $x$  is price of the car in L.E and  $n$  is the time in years from the moment of buying it . Find:
- a car price when it was brand new.      b car price after 3 years of its buying date.



## Exercises 2 - 2



- 1 Graph each of the following functions, then find the domain and range of each and show which of them is increasing and which is decreasing.
- a  $f(x) = 2^x$       b  $f(x) = 3^x$       c  $f(x) = \left(\frac{1}{2}\right)^x$       d  $f(x) = 2^{-x+1}$
- 2 **Compete:**
- a the function  $f: f(x) = 2^x$  intersects Y- axis at point \_\_\_\_\_
- b the function  $f: f(x) = 2^{1-x}$  intersects Y -axis at point \_\_\_\_\_
- c if the curve of the function  $f: f(x) = a^x$  passes through point (1, 3), then  $a =$  \_\_\_\_\_
- d the curve of the function  $f: f(x) = 3^x$  is the image of the curve of the function  $g: g(x) = \left(\frac{1}{3}\right)^x$  by reflection in \_\_\_\_\_
- e the function  $f$  where  $f(x) = a^x$  is decreasing if  $a \in$  \_\_\_\_\_
- f the function  $f$  where  $f(x) = (2a)^x$  is increasing when  $a \in$  \_\_\_\_\_
- 3 **Population:** if the population of a county by the end of year 2000 was 43265341 populations and the rate of population increasing is 1.5% yearly:
- a find a formula to represent the populations of this country after  $n$  year from year 2000..
- b use this formula to find the populations expected in year 2020 in case the increasing remains at the same rate.
- 4 **Investment:** if a man has invested 1000000LE in a project in a way that this a mount of money grows according to an exponential function with yearly increase of 6%, find:
- a A formula showing the growth of this money after  $n$  year.
- b Estimate this money after 10 years.
- 5 Find the sum of 8000LE deposited in a bank giving a yearly compound profit of 5 % for 7 years.
- 6 **Fish wealth:** if the number of Salmons in a lake is increasing according to the function of the exponential growth  $f: f(n) = 200 (1.03)^n$ , where  $n$  is the number of weeks, find the number of Salmons in this lake after 8 weeks.
- 7 If  $f(x) = 5^{x-1}$ , Prove that  $\frac{f(x) \times f(x-1)}{f(x-2) \times f(x+1)} = 1$

**You will learn**

- Power function.
- Representing the power functions graphically.
- Properties of power function.

**key terms**


- Power function
- Graphical solution

**Materials**

- Scientific calculator
- Graphic programs

**Think and discuss**

Amiba reproduce by binary fission where one cell is divided into two cells after a certain period of time, then every new cell is divided into two other cells at the same period of time in the same conditions and so on.



- Find the number of cells resulted from one cell after nine periods of time.
- Find the number of periods of time required to produce 8192 cells out of this cell.

**Learn****Exponential function**

if the equation includes one variable in the exponent, it's called an exponential function such as  $(2^{x+1} = 8)$ .

**Solving power Equation :**

**First:** if  $a^m = a^n$  where  $a \notin \{0, 1, -1\}$ , then  $m = n$ .

**Example**

- Find the solution set for each of the following equations in R:

**a**  $2^{x+3} = 8$

**b**  $3^{x-2} = \left(\frac{1}{27}\right)^x$

**Solution**

**a**  $\therefore 2^{x+3} = 8$

$\therefore 2^{x+3} = 2^3$

$\therefore x + 3 = 3$

thus,  $x = \text{zero}$

$\therefore$  Solution set = {zero}

**b**  $\therefore 3^{x-2} = \left(\frac{1}{27}\right)^x$

$\therefore 3^{x-2} = 3^{-3x}$

$\therefore x - 2 = -3x$

$\therefore x + 3x = 2$

$\therefore 4x = 2$

thus  $x = \frac{1}{2}$

$\therefore$  Solution set =  $\left\{\frac{1}{2}\right\}$

**P Try to solve**

① Find the solution set for each of the following equations in  $\mathbb{R}$ :

a  $5^{x+1} = 25$

b  $2^{1-x^2} = \frac{1}{8}$

second: if  $a^m = b^m$  where  $a, b \notin \{0, 1, -1\}$ ,

either: 1-  $m = \text{zero}$

or: 2-  $a = b$  when  $m$  is an odd number.

3-  $a = \pm b$  when  $m$  is an even number.

**Example**

② Find the solution set for each of the following equations in  $\mathbb{R}$ :

a  $3^{x+2} = 7^{x+2}$

b  $4^{x-2} = 3^{2x-4}$

**Solution**

a  $\because 3^{x+2} = 7^{x+2}$

$\therefore x + 2 = \text{zero}$  thus  $x = -2$

$\therefore$  Solution set =  $\{-2\}$

b  $\because 4^{x-2} = 3^{2x-4}$   $\therefore 4^{x-2} = 3^{2(x-2)}$

$\therefore 4^{x-2} = 9^{x-2}$

$\therefore x - 2 = \text{zero}$  thus  $x = 2$

$\therefore$  Solution set =  $\{2\}$

**P Try to solve**

② Find the solution set for each of the following equations in  $\mathbb{R}$ :

a  $5^{x-1} = 4^{x-1}$

b  $2^{2x-6} = 7^{x-3}$

**Example**

③ If  $f(x) = 2^{x+1}$ , find the value of  $x$  which satisfies  $f(x) = 32$

**Solution**

$\because f(x) = 32$   $\therefore 2^{x+1} = 32$

$\therefore 2^{x+1} = 2^5$   $\therefore x + 1 = 5$

$\therefore x = 4$   $\therefore$  Solution set =  $\{4\}$

**P Try to solve**

③ If  $f(x) = 7^x$ , find the value of  $x$  which satisfies  $f(x+1) = 49$

## Solving the power Equations Graphically

 **Example**

- 4 in one figure, graph the curve of each of the two functions  $f_1$  where  $f_1(x) = 2^x$  and  $f_2$  where  $f_2(x) = 6 - x$ , then find the solution set of the equation  $2^x = 6 - x$  from the graph.

 **Solution**

$x$	-3	-2	-1	0	1	2	3
$2^x$	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	4	8
$6 - x$	9	8	7	6	5	4	3

from the graph, the  $x$  coordinate of the intersection point equals 2  
 $\therefore$  the solution set of the equation =  $\{2\}$


 **Try to solve**

- 4 use one of the graphic programs (geogebra) to graph each of the two functions  $f_1(x) = 2^{x+1}$  and  $f_2(x) = 3$  in one figure. From the graph, find the solution set of the equation  $2^{x+1} = 3$ .

 **Example**

- 5 **Biology:** A microorganism reproduces by binary fission where the number of these organisms is replicated each hour because each cell is divided into two cells. If the number of cells at the beginning was 20000 cells, find:

- Number of cells after 5 hours.
- After how many hours does the number of cells get 2560000?

 **Solution**

the number of cells can be written in the form of an exponential function.

$$\begin{aligned} f(n) &= b(a)^n \\ &= 20000(2)^n \quad \text{where } n \text{ is number of hours} \end{aligned}$$

- Number of cells after 5 hours (by placing  $n = 5$ )  
 $= 20000 \times 2^5 = 640000$  cells
- To find how many hours required to get 2560000, place  $f(x) = 2560000$   
 $\therefore 20000(2)^n = 2560000$  **By dividing by 20000**  
 $\therefore 2^n = 128$   
 $\therefore 2^n = 2^7$  **thus  $n = 7$  hours.**

 **Try to solve**

- 5 Answer the questions mentioned in **think and discuss** page (66)





## Exercises 2 - 3



## 1 Complete:

- a if  $5^{x-2} = 1$  then  $x =$  \_\_\_\_\_
- b if  $3^{x-2} = 7^{x-2}$  then  $x =$  \_\_\_\_\_
- c if  $2^{x+1} = 5^{x+1}$  then  $3^{x+1} =$  \_\_\_\_\_
- d if  $2^{|x|} = 32$  then  $x =$  \_\_\_\_\_
- e If the curve of the function  $f_1$  where  $f_1(x) = 3^x$  intersects the curve of the function  $f_2$  where  $f_2(x) = 4-x$  at point  $(k, 3)$ , then the solution set of the equation  $3^x = 4-x$  is \_\_\_\_\_

## Choose the correct answer

- 2 if  $3^{x-5} = 9$ , then  $x =$  \_\_\_\_\_
- a 2                      b 7                      c -3                      d -7
- 3 if  $2^x = 20$  where  $n < x < n + 1$  and  $n$  is an integer, then  $n =$  \_\_\_\_\_
- a 1                      b 2                      c 3                      d 4
- 4 if  $3^x = 9$ , then  $3^{x+1} =$  \_\_\_\_\_
- a 5                      b 15                      c 27                      d 45
- 5 The number  $5^{x+1} + 5^x$  is divisible by \_\_\_\_\_ for all the natural values of  $x$ .
- a 7                      b 6                      c 13                      d 17
- 6 if  $(\frac{2}{3})^{x-2} = \frac{8}{28}$ , then  $x =$  \_\_\_\_\_
- a 2                      b 3                      c 4                      d 5
- 7 The two curves of the two functions  $f(x) = 2^x$  and  $g(x) = 3^x$  intersect at  $x =$  \_\_\_\_\_
- a -1                      b 0                      c 1                      d 2
- 8 Find the solution set for each of the following equations in  $\mathbb{R}$ :
- a  $3^{x+4} = 9$                       b  $2^{x-5} = \frac{1}{32}$
- c  $5^{x+2} = 1$                       d  $3^{|x|} = 3$

**e**  $2 \times 3^{x-2} = 54$                       **f**  $7^{x-5} = 3^{x-5}$   
**g**  $2^{3x-6} = 5^{x-2}$                       **h**  $(\frac{3}{2})^{x-2} = \frac{8}{27}$   
**i**  $2^x \times 5^{-x} = \frac{4}{25}$                       **j**  $4^x = 64$   
**k**  $4^{1-x} = \frac{1}{4}$                                   **l**  $(3)^{x-5} = \frac{1}{9}$

9 Find the solution set for each of the following equations graphically:

**a**  $3^x = 3$                                   **b**  $2^{x+1} = 5$       Rounding the sum to one decimal  
**c**  $3^{x+1} = -x$                           **d**  $2x = \frac{1}{2}x + 1$

10 If  $f(x) = 2^x$ , find the solution set for each of the following equations:

**a**  $f(x) = 8$                               **b**  $f(x+1) = \frac{1}{32}$

11 If  $f(x) = 3^{x+1}$ , find the solution set for each of the following equations:

**a**  $f(x) = 27$                               **b**  $f(x-1) = \frac{1}{9}$

12 If  $f(x) = 7^{x-2}$ , find the solution set for each of the following equations:

**a**  $f(x) = 343$                               **b**  $f(2x) = \frac{1}{49}$

13 **Discover the error:** Mohammed and Karem have solved the equation  $2 \times 2^x = 16$  as follows:

**Mohammed's Solution**

$2 \times 2^x = 16$   
 $\therefore 4^x = 16$   
 $\therefore 4^x = 4^2$   
 $\therefore x = 2$

**Karem's solution**

$2 \times 2^x = 16$   
 $\therefore 2^x = \frac{16}{2} = 8$   
 $\therefore 2^x = 2^3$   
 $\therefore x = 3$

What is the right solution? Why?

14 the number of marine organisms decreases according to the function of the exponential decay  $y = 8192 (\frac{1}{2})^{n-1}$  where  $n$  is the number of weeks from now. Find:

- a** The number of these organisms after 4 weeks from now.  
**b** After how many weeks does the number of these organism get 256?

# Logarithmic Function and its Graphical Representation

## 2 - 4



### Think and discuss

Check the following exponential functions and try to solve each of them: if  $2^x = 2$ ,  $2^y = 4$ ,  $2^z = 3$ , then:

1-  $x = \underline{\hspace{2cm}}$ ,  $y = \underline{\hspace{2cm}}$

2- The value of  $z$  is included between two consecutive integers which are  $\underline{\hspace{2cm}}$  and  $\underline{\hspace{2cm}}$

**Notice that** the value of  $y$  cannot be calculated directly such as  $x$  and  $z$ , so we need to the concept of a new function to calculated the value of  $y$ .



### Learn

### Logarithmic Function

if  $x$  and  $a$  are two positive numbers where  $a \neq 1$ , then the logarithmic function  $y = \log_a x$  is the inverse function of the power function  $y = a^x$

**Example:** if  $\log_2 32 = 5$ , then  $2^5 = 32$  and vice versa.

#### Verbal expression:

If point  $(c, d) \in$  exponential function  $y = a^x$ , then:

1- point  $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}) \in$  function  $y = \log_a x$ .

2- The exponential form  $a^c = d$  where  $a \in \mathbb{R}^+ - \{1\}$  is equivalent to the logarithmic form  $\underline{\hspace{2cm}}$



### Example

#### Converting into the logarithmic form.

① convert each of the following into the logarithmic form:

a  $3^4 = 81$

b  $25^{\frac{1}{2}} = \frac{1}{5}$

c  $10^{-2} = 0.01$

#### Solution

a  $\log_3 81 = 4$

b  $\log_{25} \frac{1}{5} = -\frac{1}{2}$

c  $\log_{10} 0.01 = -2$

**Verbal expression:** Can we convert  $(-2)^4 = 16$  into a logarithmic form? Explain.

### You will learn

- Definition of the logarithmic function.
- Graphical representation of the logarithmic function.
- Converting from the exponential form into the logarithmic form and vice versa.
- Solving some simple logarithmic equation.

### Key terms

- logarithm
- inverse function
- domain
- common logarithm

### Materials

- Calculator.
- Computer.



### Tip

$\log_a x = y$  is called the logarithmic form while  $a^y = x$  is called the equivalent exponential form. Notice that (a) is a positive base. If  $(-3)^4 = 81$ , then there is not a logarithmic form equivalent to it.

**P Try to solve**


1 Express each of the following in a logarithmic form:

a  $10^3 = 1000$

b  $8^{\frac{1}{3}} = 2$

c  $b^x = y$  where  $b \in \mathbb{R}^+ - \{1\}$

**Common Logarithm**

it is the logarithm whose base is 10 and written without writing the base. I.e.  $\log_{10} 7 = \log 7$  and  $\log_{10} 127 = \log 127$ . The button  in the calculator can be used to find the common logarithm of any number.

**Example**

2 convert each of the following into the exponential form:

a  $\log_2 32 = 5$

b  $\log 1000 = 3$

c  $\log_2 1 = \text{zero}$

**Solution**

a  $2^5 = 32$

b  $10^3 = 1000$

c  $2^{\text{zero}} = 1$

**P Try to solve**

2 convert each of the following into the exponential form:

a  $\log_{125} 25 = \frac{2}{3}$

b  $\log 100 = 2$

c  $\log_5 5 = 1$

**Example**

**Finding the values of logarithmic phrases**

3 Find the value of each:

a  $\log_5 125$

b  $\log 0.01$

**Solution**

a Let  $\log_5 125 = x$  and by converting into the exponential form:

$\therefore 5^x = 125$                        $\therefore 5^x = 5^3$                       thus  $x = 3$

$\therefore \log_5 125 = 3$

b Let  $\text{Log}0.01 = y$  (a common logarithm whose base is 10) and by converting into the exponential form.

$\therefore 10^y = 0.01$                        $\therefore 10^y = 10^{-2}$

thus  $y = -2$                        $\therefore \log 0.01 = -2$

**Try to solve**

3 Find the value of each:

a  $\log_3 81$

b  $\log_{\frac{1}{4}} 32$

**Example Solving the equations**

4 Find the solution set for each of the following equations in R:

a  $\log_2 (x+5) = 3$

b  $\log_5 625 = x - 1$

c  $\log_x (x+6) = 2$

**Solution**

a The equation is defined for all the values of  $x + 5 > \text{zero}$  i.e.  $x > -5$  (is the domain of defining the equation).

By converting the equation into the exponential form

$\therefore x + 5 = 2^3$   $\therefore x + 5 = 8$

thus  $x = 3$

$\therefore 3 \in$  the domain of defining the equation  $\therefore$  Solution set = **{3}**

b The equation is defined for all real values of  $x$  and by converting the equation into the exponential form

$\therefore 5^{x-1} = 625$   $\therefore 5^{x-1} = 5^4$

$\therefore x - 1 = 4$  thus  $x = 5$

$\therefore$  Solution set = **{5}**

c The equation is defined for all the values of  $x$  which satisfy each of  $\begin{cases} x + 6 > \text{zero} \\ x > \text{zero} \\ x \neq 1 \end{cases}$

i.e. the domain of defining the equation is  $[\text{zero}, \infty] - \{1\}$

and by converting the equation into the exponential form:

$x^2 = x + 6$   $x^2 - x - 6 = 0$

$(x - 3)(x + 2) = 0$

either  $x = 3$  or  $x = -2$

Since  $x = -2 \notin$  the domain of defining the equation

$\therefore$  Solution set = **{3}**

**Try to solve**

4 Find the solution set for each of the following equations in R:

a  $\log_5 (3x-1) = 1$

b  $\log_3 27 = x + 2$


c  $\log_{(x-1)} 9 = 2$

### Graphical Representation of the Logarithmic Function

If  $f(x) = a^x$  where  $a \in \mathbb{R}^+ - \{1\}$ , then the inverse function of the function  $f$  is called the logarithmic function. I.e.  $y = \log_a x$

### Relation between the exponential and logarithmic function

The opposite figure represents the exponential function  $y = a^x$  and logarithmic function  $y = \log_a x$ . Study the properties of both functions for domain, range, monotony and symmetry around the straight line  $y = x$  in one figure.




#### Example

- 1 Graph the curve of each of the two functions  $y = \log_2 x$  and  $y = \log_{\frac{1}{2}} x$  in one figure.

#### Solution

we choose values of  $x$  the powers of the number 2 ( the base)  $\{2^{-2}, 2^{-1}, 2^0, 2^1, 2^2\}$

$x$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	4
$\log_2 x$	-2	-1	zero	1	2
$\log_{\frac{1}{2}} x$	2	1	zero	-1	-2



From the graph, you can deduce the following properties for the curve of the logarithmic function:

the domain =  $\mathbb{R}^+$ , and the range =  $\mathbb{R}$

the function  $y = \log_a x$  is increasing for each  $a > 1$  and is decreasing for each  $0 < a < 1$

#### Try to solve

- 5 Represent the curve of the function  $y = \log_3 x$  graphically, then find the range of the function and investigate its monotony from the graph.

#### Example

- 2 **Life applications:** A country applies a tax system where the financier pays the yearly dues according to the country's system.

$$f(x) = \begin{cases} 10\% \cdot x & \text{when } x \leq 5000 \\ 10\% \cdot x + 100 \log(x - 4999), & \text{when } x > 5000 \end{cases}$$

where  $x$  is the yearly net profit, **find:**

- The tax dues that a financier whose yearly net profit is 3600 L.E.
- The tax dues that a financier whose yearly net profit is 8000 L.E.

**Solution**

- a  $f(3600) = 10\% \times 3600 = 0.1 \times 3600 = 360$  L.E  
 b  $f(8000) = 10\% \times 8000 + 100 \log (8000 - 4999) = 1147.7$  L.E

**Try to solve**

- 6 If  $a$  expresses the amount of money spent for an advertisement of a company in the year and  $y$  expresses the amount of money which the company gets after the sales of this year where  $y = 10^4 [1 + 2 \log (\frac{a}{100} + 1)]$ . Calculate  $y$  when  $a = 1100$  L.E.


**Exercise 2 - 4**

**1 Complete:**

- a The exponential form equivalent to the form  $\log_3 27 = 3$  is \_\_\_\_\_  
 b The logarithmic form equivalent to the form  $3^{\text{zero}} = 1$  is \_\_\_\_\_  
 c  $\log 0.001 =$  \_\_\_\_\_ d  $\log_2 1 =$  \_\_\_\_\_  
 e If  $\log_x 4 = 2$ , then  $x =$  \_\_\_\_\_ f if  $\log_2 128 = x + 1$ , then  $x =$  \_\_\_\_\_  
 g The domain of the function  $f: f(x) = \log_2 x$  is \_\_\_\_\_  
 h The function  $f$  where  $f(x) = \log_a x$  is decreasing for each  $a \in$  \_\_\_\_  
 i The curve of the function  $f$  where  $f(x) = \log_2 x$  passes through point  $(8, \text{_____})$   
 j If  $\text{Log} 3 = x$  and  $\text{Log} 5 = y$ , then  $\log 15 =$  \_\_\_\_\_ (in terms of  $x, y$ )
- 2 Find the solution set for each of the following equations in  $\mathbb{R}$ :
- a  $\log_3 (x - 1) = 2$       b  $\log_5 (x + 2) = 3$       c  $\log_x 9 = \frac{2}{3}$   
 d  $\log_{x+1} 8 = \frac{3}{4}$       e  $\log_x (x + 2) = 2$       f  $\log_x 9 = 2$
- 3 Find the value of the following without using the calculator.
- a  $\log_5 1$       b  $\log_7 7$       c  $\log_3 9$       d  $\log_6 3 + \log_6 2$
- 4 Represent the function  $f$  in each of the following graphically, then find its range and investigate its monotony:
- a  $f(x) = \log_2 x$       b  $f(x) = \log_3 x$       c  $f(x) = \log_{\frac{1}{2}} x$       d  $f(x) = \log_{\frac{1}{3}} (x + 1)$
- 5 use the calculator to find the value of the following:
- a  $\log 15$       b  $\log_2 27$       c  $4 \log 7 - 5 \log 13$
- 6 If the cost of the annual subscription of a family in a social club in pounds follows the relation  $f(x) = 500 + 100 \log (n x)$ , where  $n$  is the number of subscription years and  $x$  is the family member. Find the value of subscription of a five -member family for the fourth year in this club.

## Some Properties of Logarithms

**You will learn**

- using some properties of logarithms.
- Solving logarithmic equations.
- Using the calculator to solve the exponential functions.
- Life applications on logarithms.

**Key terms**

- logarithmic equation

**Materials**

- Scientific calculator
- Computer with graphic programs

In the previous lesson, you learned the concept of a logarithm and how to represent the logarithmic function graphically. Now, we are going to list some properties of logarithms to help you simplify the logarithmic expressions or solve the equations containing a logarithm.

**Learn****Some Properties of Logarithms**

if  $a \in \mathbb{R}^+ - \{1\}$ ,  $x, y \in \mathbb{R}^+$ , then

$$1- \log_a a = 1$$

For example,  $\log_3 3 = 1$ ,  $\log_{10} 10 = 1$

$$2- \log_a 1 = \text{zero}$$

For example,  $\log_5 1 = \text{zero}$ ,  $\log 1 = \text{zero}$

Try to prove **1** and **2** from the definition of logarithms

**3- Multiplication property in logarithms:**

$$\log_a xy = \log_a x + \log_a y \quad \text{where } x \text{ and } y \in \mathbb{R}^+$$

To prove the correctness of this property:

$$\text{place } b = \log_a x \text{ and } c = \log_a y$$

From the definition of logarithms, we find that:

$$x = a^b, y = a^c$$

$$\text{then } xy = a^b \times a^c \quad \text{Le } xy = a^{b+c}$$

By converting this form into the logarithmic form, then:

$$\log_a xy = b + c$$

By substituting the two values of  $b$  and  $c$ , then  $\log_a xy = \log_a x + \log_a y$

**Example**

- ① Find the value of  $\log_{34} 2 + \log_{34} 17$  without using the calculator.



**Solution**

$$\begin{aligned} \text{the expression} &= \log_{34} (2 \times 17) && \text{use property (3)} \\ &= \log_{34} 34 \\ &= 1 && \text{use property (1)} \end{aligned}$$

**Try to solve**

- 1 if  $\log_2 7 \approx 2.8$ ,  $\log_2 13 \approx 3.7$ , find the value of  $\log_2 91$  without using the calculator.

**4- Division property in logarithms:**

$$\log_a \frac{x}{y} = \log_a x - \log_a y \quad (\text{try to prove the correctness of this relation yourself})$$

**Example**

- 2 Find the value of  $\log 50 - \log 5$  without using the calculator.

**Solution**

$$\begin{aligned} \text{the expression} &= \log \frac{50}{5} && \text{use the division property} \\ &= \log 10 = 1 && \text{use property (1)} \end{aligned}$$

**Try to solve**

- 2 Find the value of  $\log_2 7 - \log_2 3.5$  without using the calculator.

**5- Property of power logarithm:**

$$\log_a x^n = n \log_a x \quad \text{where } x > 0 \quad (\text{try to prove the correctness of the relation yourself})$$

**Example**

- 3 Find the value of  $\log_5 125$  without using the calculator.

**Solution**

$$\begin{aligned} \text{the expression} &= \log_5 35 \\ &= 3 \log_5 5 && \text{use the property of power} \\ &= 3 \times 1 = 3 && \text{use property (1)} \end{aligned}$$

Notice that:  $\log_a \left(\frac{1}{x}\right) = -\log_a x$  where  $x \in \mathbb{R}^+$

**Try to solve**

- 3 Find  $\log_{\sqrt{3}} 27$  in the simplest form.
- 4 **Critical thinking:** is the domain of the function  $f(x) = \log_a x^2$  the same as the domain of the function  $g(x) = 2 \log_a x$ ? Explain.

## 6 - Property of changing the base

$$\log_y x = \frac{\log_a x}{\log_a y}$$

and proving the correctness of this property

By placing:  $z = \log_y x$ 


$$y^z = x$$

by converting into the exponential form

$$z \log_a y = \log_a x$$

the logarithm of the two sides is taken off for the base a

$$\text{then } z = \frac{\log_a x}{\log_a y} \quad \text{I.e: } \log_y x = \frac{\log_a x}{\log_a y}$$

 Example

- ④ Reduce to the simplest form  $\log_7 16 \times \log_2 49$

 Solution

$$\begin{aligned} \text{the expression} &= \frac{\log 16}{\log 7} \times \frac{\log 49}{\log 2} && \text{use property (6)} \\ &= \frac{\log 2^4}{\log 7} \times \frac{\log 7^2}{\log 2} \\ &= \frac{4 \log 2}{\log 7} \times \frac{2 \log 7}{\log 2} && \text{use property (5)} \\ &= 4 \times 2 = 8 \end{aligned}$$

 Try to solve

- ⑤ Find the solution of the example above by changing the base into another number but not 10.

## 7 - Property of the multiplicative inverse.

$$\log_b a = \frac{1}{\log_a b} \quad \text{I.e both } \log_b a \text{ and } \log_a b \text{ are multiplicative inverse of each other (try to}$$

prove the correctness of this relation ).

 Example

- ⑤ Find the value of  $\frac{1}{\log_3 15} + \frac{1}{\log_5 15}$  without using the calculator.

 Solution

$$\begin{aligned} \text{the expression} &= \log_{15} 3 + \log_{15} 5 && \text{use property (7)} \\ &= \log_{15} (3 \times 5) && \text{use property (3)} \\ &= \log_{15} 15 = 1 && \text{use property (1)} \end{aligned}$$

**Try to solve**

- 6 Find the value of  $\frac{1}{\log_2 30} + \frac{1}{\log_3 30} + \frac{1}{\log_5 30}$  without using the calculator.

**Simplifying the Logarithmic Expressions****Example**

- 6 Reduce  $\log 0.009 - \log \frac{27}{16} + 3 \log \frac{5}{2} - \log \frac{1}{12}$  to the simplest form

**Solution**

$$\begin{aligned} \text{the expression} &= \log \frac{9}{1000} - \log \frac{27}{16} + \log \left(\frac{5}{2}\right)^3 - \log \frac{1}{12} && \text{property (5)} \\ &= \log \left(\frac{9}{1000} \times \frac{16}{27} \times \frac{125}{8} \times \frac{12}{1}\right) && \text{property (3), (4)} \\ &= \log 1 = \text{zero} && \text{property (2)} \end{aligned}$$

**Try to solve**

- 7 Reduce  $4 \log \sqrt{3} - \log_5^2 1 - \log_7^2 1 - \log \frac{1}{2}$  to the simplest form.

**Solving Logarithmic Equations****Example**

- 7 Find the solution set for each of the following equations in R.

a  $\log_2 x + \log_2 (x+1) = 1$

b  $\log_2 x + \log_4 x = 3$

**Solution**

- a The function is defined for each  $x > \text{zero}$  and  $x+1 > \text{zero}$

i.e.  $x > \text{zero}$  (domain of defining the equation)

$$\therefore \log_2 x(x+1) = 1 \quad \text{use property (3)}$$

$$\therefore x(x+1) = 2^1 \quad \text{converting from the logarithmic form into the exponential form}$$

$$\therefore x^2 + x - 2 = \text{zero} \quad \therefore (x+2)(x-1) = \text{zero}$$

Either  $x = -2$  or  $x = 1$ , and  $x = -2 \notin$  the domain of defining the equation

$$\therefore \text{Solution set} = \{1\}$$

- b The function is defined for each  $x > \text{zero}$  (domain of defining the equation)

$$\therefore \log_2 x + \frac{\log_2 x}{\log_2 4} = 3 \quad \text{property (6)}$$

$$\log_2 x + \frac{\log_2 x}{2} = 3 \quad \text{multiply by 2}$$

$$\therefore 2 \log_2 x + \log_2 x = 6 \quad \therefore 3 \log_2 x = 6 \quad \therefore \log_2 x = 2$$

$$\therefore x = 4 \quad \text{(converting from the logarithmic form into the exponential form)}$$

where  $x = 4 \in$  the domain of defining the equation

$$\therefore \text{Solution set} = \{4\}$$

**P Try to solve**

 8 Find the solution set for each of the following equations in  $\mathbb{R}$ :

a  $\log(2x + 1) - \log(3x - 1) = 1$

b  $\log_2 x = \log_x 2$

**Solving the Exponential functions by Using Logarithms**
**Example**

 9 Find the solution set for each of the following equations in  $\mathbb{R}$  rounding the sum to the nearest two decimals :

a  $2x = 7$

b  $3 \cdot x^{+1} = 5x^{-2}$

**Solution**

a  $2x = 7$

take off the logarithm of two sides

$$\therefore \log 2^x = \log 7$$

$$\therefore x \log 2 = \log 7$$

$$\therefore x = \frac{\log 7}{\log 2}$$

use the calculator respectively as follows:

$$\log 7 \div \log 2 = 2.807354922$$

$$\therefore x \approx 2.81$$

$$\therefore \text{The solution set} = \{2.81\}$$

(check your answer using the calculator)

$$2 \cdot x^{\wedge} \text{ans} = 7$$

b  $3^{x+1} = 5^{x-2}$

take of the log of two sides

$$\therefore (x+1)\log 3 = (x-2)\log 5$$

$$\therefore x \log 3 + \log 3 = x \log 5 - 2 \log 5$$

$$\therefore x \log 3 - x \log 5 = -\log 3 - 2 \log 5$$

$$\therefore x(\log 3 - \log 5) = -\log 3 - 2 \log 5$$

$$\therefore x = \frac{-\log 3 - 2 \log 5}{\log 3 - \log 5}$$

use the calculator respectively as follows:

$$\log 3 \div (\log 3 - \log 5) = 2.807354922$$

$$\therefore x \approx 8.45$$

$$\therefore \text{The solution set} = \{8.45\}$$

(check your answer using the calculator)

$$3 \cdot x^{\wedge} \text{ans} + 1 = 5 \cdot x^{\wedge} \text{ans} - 2$$

**P Try to solve**

8 Find the solution set of each of the equations to the nearest two decimals:

a  $7^x = 2$

b  $4^{x-1} = 3^x$


**Activity**
**Mathematical and life applications**

 10 **Industry** : If the working efficiency of a machine decreases annually according to the relation  $E = (0.9)^n$  where  $E$  is the machine efficiency,  $E_0$  is the machine initial efficiency, and  $n$  is the number of working years of the machine. If it is known that the machine is no longer used to work if its efficiency reaches 40%. How many years can this machine work before it stops working?

**Solution**

It's meant by reaching the efficiency 40% that 40% is calculated from the initial efficiency.

$$\begin{aligned} \therefore 0.4 E_0 &= (0.9)^n E_0 && \text{divide by } E_0 \\ \therefore 0.4 &= (0.9)^n && \text{take off the log of two sides} \\ \therefore \log 0.4 &= n \log 0.9 && \therefore n = \frac{\log 0.4}{\log 0.9} = 8.696718 \end{aligned}$$

I.e. the machine works more than 8 years and half a year.

**P Application on the activity**

9 In the example above, find the machine efficiency after 4 years of its working date .


**Exercises 2 - 5**


choose the correct answer:

- 1  $\log_2 8 =$ 
  - a 4
  - b 3
  - c 16
  - d 10
- 2  $\text{Log}2 + \text{Log}5 =$ 
  - a 1
  - b  $\log 7$
  - c  $\text{Log}2.5$
  - d 10
- 3  $\log_5 \sqrt{5} =$ 
  - a 2
  - b 5
  - c  $\frac{1}{2}$
  - d -1
- 4 if  $\log 3 = x$  ,  $\log 4 = y$  , then  $\log 12 =$ 
  - a  $x + y$
  - b  $xy$
  - c  $x - y$
  - d  $\log x + \log y$
- 5  $2 \log_6 2 + 2 \log_6 3 =$ 
  - a 6
  - b 36
  - c 2
  - d 12
- 6  $\log_2 5 \times \log_5 2 =$ 
  - a 1
  - b 10
  - c  $\frac{5}{2}$
  - d zero
- 7  $\log_5 2 \times \log_3 5 \times \log_2 3 =$ 
  - a 30
  - b 1
  - c zero
  - d  $\log 30$
- 8 Express each of the following in terms of  $\log x$  and  $\log (x + 1)$ 
  - a  $\log x (x + 1)$
  - b  $\log \frac{x}{x + 1}$
  - c  $\log \sqrt{x} (x + 1)^2$
- 9 Reduce to the simplest form:
  - a  $\log_6 54 - \log_6 9$
  - b  $\log_6 2 + \log_6 3$
  - c  $\log_2 12 + \log_2 \frac{2}{3}$
  - d  $\log 48 + \text{Log}125 - \log 6$
  - e  $\frac{1 - \text{Log}2}{\text{Log}125}$
  - f  $\frac{\text{Log}49 + 3 \log 7}{\log 7}$
  - g  $\log_2 16 + \log_3 \sqrt{3} + \log 0.1$

h  $\frac{1}{2} \log_3 a + \frac{1}{2} \log_3 b + 2 \log_3 c - \log_3 \sqrt{ab} - \log_3 3c^2$

10 Find the solution set for each of the following equations in  $\mathbb{R}$ :

a  $\log_2 x + \log_2 (x+2) = 3$

b  $\log x + \log (x-3) = 1$

c  $\log_5 x - \log_5 2 = 2$

d  $\log (x+3) - \log 3 = \log x$

e  $\frac{1}{\log_2 x} + \frac{1}{\log_3 x} = 2$

f  $\log x - \frac{3}{\text{Log} x} = 2$

11 Prove that  $\log_b a \times \log_c b \times \log_d c \times \log_a d = 1$ , then calculate the value of

$\log_2 3 \times \log_3 5 \times \log_5 16$

12 Find the value of  $x$  in each of the following, then round the sum to the nearest decimal.

a  $3^x = 7$

b  $5^{x-1} = 2$

c  $4 \times 7^{x-2} = 1$

d  $2^{x-3} = 3^{x+1}$



### Activity

**Biology:** If the volume of a sample of bacteria in a certain moment is  $3 \times 10^6$  and the volume of the sample increases according to the power function  $v = 3 \times 10^6 (1.15)^n$ , find the volume of bacteria after 4 hours.

### @ Enrichment Information

Please visit the following link.



## Unit summary

## 1) The integer exponents

- a**  $a^n = a \times a \times a \times \dots \times a$  (factor  $a$  is repeated  $n$  times)  
**b**  $a^0 = 1$  where  $a \in \mathbb{R} \setminus \{0\}$  } **c**  $a^{-1} = \frac{1}{a}$ ,  $a^{-n} = \frac{1}{a^n}$  where  $a \neq 0$

## Properties of integer exponents

for each  $m, n \in \mathbb{Z}$  and  $a, b \in \mathbb{R}$ ,  $b \neq 0$ , then:

- a**  $a^m \times a^n = a^{m+n}$       **b**  $(ab)^n = a^n b^n$       **c**  $\frac{a^m}{a^n} = a^{m-n}$   
**d**  $\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$       **e**  $(a^m)^n = a^{mn}$

2) The  $n^{\text{th}}$  roots:

The equation  $x^n = a$  where  $a \in \mathbb{R}$ ,  $n \in \mathbb{Z}^+$  has  $n$  of roots

- a**  $n$  is an even number,  $a \in \mathbb{R}^+$   
 There are two real roots (the rest of the roots are complex numbers) one root is positive while the other is negative and both are denoted by  $\pm \sqrt[n]{a}$ .  
**b**  $n$  is an even number,  $a \in \mathbb{R}^-$ , the equation doesn't have real roots (all the roots are complex number).  
**c**  $n$  is an odd number,  $a \in \mathbb{R}$ , the equation has only one real root (the rest of the roots are complex numbers) and is denoted by  $\sqrt[n]{a}$ .  
**d**  $n \in \mathbb{Z}^+$ ,  $a = \text{zero}$   
 The equation has only one solution which is zero (it has  $n$  of repeated roots and each root equal zero)

3) Properties of  $n^{\text{th}}$  roots:

if  $\sqrt[n]{a}$  and  $\sqrt[n]{b} \in \mathbb{R}$ , then:

- a**  $\sqrt[n]{ab} = \sqrt[n]{a} \times \sqrt[n]{b}$       **b**  $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$ ,  $b \neq 0$   
**c**  $\sqrt[n]{a^m} = (\sqrt[n]{a})^m = a^{\frac{m}{n}}$       **d**  $\sqrt[n]{a^n} = a$  if  $n$  is an odd number and equals  $|a|$  if  $n$  is an even number

4) Rational exponents  $a^{\frac{1}{n}} = \sqrt[n]{a}$ 

## 5) Properties of Rational exponents

- a**  $a^{\frac{m}{n}} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m \in \mathbb{R}$  where  $a \in \mathbb{R}$ ,  $n \in \mathbb{Z}_+ - \{1\}$ ,  $m \in \mathbb{Z}$  and there is no common factor between  $m$  and  $n$ .  
**b** The rules of integer powers can be generalized on rational powers.

6) The power function: if  $f: \mathbb{R} \rightarrow \mathbb{R}^+$  where  $f(x) = a^x$  for each  $a \in \mathbb{R}^+ - \{1\}$ , then  $f$  is called a power function whose base is  $a$ .

**7 Properties of the curve of The power function**

- a** The domain of the function =  $\mathbb{R}$
- b** the range  $\mathbb{R}^+$
- c** The function is increasing on its domain for each  $a > 1$  and called the function of the exponential growth.
- d** The function is decreasing on its domain for each  $0 < a < 1$  and called the function of the exponential decay.

**8 Exponential growth:** The function  $f$  where  $f(x) = a(1 + g)^n$  can be used to represent the exponential growth by a constant percentage in equal periods of time where  $n$  is the period of time,  $a$  is the initial value and  $g$  is the percentage of growth in one period of time

**9 Exponential decay:** The function  $f$  where  $f(x) = a(1 - g)^n$  can be used to represent the exponential decay by a constant percentage in equal periods of time where  $t$  is the period of time,  $a$  is the initial value and  $g$  is the percentage of decay in one period of time.

**10 Logarithmic function**

- a** if  $a \in \mathbb{R}^+ - \{1\}$ , then the function  $y = \log_a x$  is an inverse function to the power function  $y = a^x$
- b**  $a^b = c$ , then  $b = \log_a c$  (converting from the exponential form into the logarithmic form and vice versa).
- c** The common logarithm is logarithm whose base is 10 {notice that  $\log_5 = \log_{10} 5$ }

**11 Properties of the logarithmic function**

- a** The domain of the function =  $\mathbb{R}^+$
- b** The range =  $\mathbb{R}$
- c** The function  $y = \log_a x$  is increasing for each  $a > 1$  and decreasing for each  $0 < a < 1$

**12 Properties of logarithms : if  $a \in \mathbb{R}^+ - \{1\}$**

- a**  $\log_a a = 1$
- b**  $\log_a a = \text{zero}$
- c**  $\log_a x^m = m \log_a x$  where  $x > \text{zero}$
- d**  $\log_a x + \log_a y = \log_a xy$  where  $x, y > \text{zero}$
- e**  $\log_a x - \log_a y = \log_a \frac{x}{y}$  where  $x, y > \text{zero}$
- f**  $\log_a x = \frac{\log_b x}{\log_b a}$  where  $x > \text{zero}$ ,  $a, b \in \mathbb{R}^+ - \{1\}$
- g**  $\log_a x \times \log_x a = 1$



**General Exercises**

For more exercises please visit the website of the Ministry of Education.





### Accumulative Test



- 1 Determine the domain of each of the following functions:
- a  $f(x) = \sqrt{x-2}$       b  $f(x) = \frac{x-2}{x}$       c  $g(x) = \log_3(x-2)$
- 2 Graph the curve of each of the following, then determine the range, investigating the monotony of the function and its type whether it is even or odd.
- a  $f(x) = (x-2)^2$       b  $f(x) = 3x-1$   
 c  $g(x) = 2 - |x|$       d  $E(x) = 1 - \log_2 x$
- 3 Reduce:
- a  $\frac{\sqrt{a} \times \sqrt[3]{a^2}}{\sqrt{a}}$       b  $(125)^{\frac{2}{3}} \times (81)^{\frac{1}{4}} \times (15)^{-1}$
- 4 Find the value of each of the following (without using the calculator).
- a  $(16)^{\frac{1}{4}}$       b  $\sqrt[3]{27^{-2}}$       c  $(3^{-1})^2 \times 9 \times (\sqrt[4]{15})^{\text{zero}}$   
 d  $\text{Log}5 + \log \frac{1}{5}$       e  $\log_{\frac{1}{9}} \frac{1}{27}$       f  $3 \log_3 5 + \log_3 \frac{243}{125}$
- 5 Find the solution set for each of the following equations in R:
- a  $|x-2| = 5$       b  $3^{x-2} = \frac{1}{3}$   
 c  $x^{-4} = \frac{1}{16}$       d  $\log x = \log 3 + \log 10$
- 6 Use the calculator to find:
- a the value of  $x$  which satisfies  $3^{x-2} = 25$  rounding the sum to the nearest two decimals.  
 b the value of  $\frac{3^{750}}{5^{310}}$
- 7 Which of the following functions represents a growth function and which represents a decay one:
- a  $y = 3(1.05)^x$       b  $y = 10(2.1)^{x+1}$   
 c  $y = 0.4(\frac{1}{2})^x$       d  $y = 0.2(3)^{1-x}$



## Unit Three

# Limits

### Unit introduction

calculus is one of the modern branches of mathematics concerning with the studying of limits, continuity, differentiation, integration and the infinite series. It is the science used to study the variation in the functions and their factorization.

Calculus deals with numerous geometrical life application, trade and different science. Calculus is basically used to study the behaviour of the function and the change in it and to solve the problems which algebra and other sciences cannot deal with.

### Unit objectives


By the end of this unit, the student should be able to:

- Identify an introduction about limits.
- Identify some unspecified quantities like:  
 $\frac{0}{0}$ ,  $\frac{\infty}{\infty}$ ,  $\infty - \infty$ ,  $0 \times \infty$
- Determine a method to find the limit of a function, by direct substitution, factorization, long division, and multiply by conjugates:
- Find the limit of a function using the rule  
 $\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = n a^{n-1}$
- Conclude the limit of a function using the rule:  
 $\lim_{x \rightarrow a} \frac{x^n - a^n}{x^m - a^m} = \frac{n}{m} a^{n-m}$
- Find the limit of a function at the infinity algebraically and graphically.
- Use the graphic calculators to check the limit of the function and the value of the limit.
- Identify various applications on the basic concepts of the function limits.

## Key terms

- Unspecified Quantity
- Undefined
- Limit of a Function
- direct Substitution
- Conjugate
- Polynomial Function
- Limit of a Function at Infinity

## Unit planning guide



## Unit Lessons

- lesson (3-1): Introduction to limits .
- lesson (3-2): Finding the limit of a function algebraically.
- lesson (3-3): Limit of a function at infinity.

## Materials

Scientific calculator - Computer - Graphical programs for PC

## Introduction to Limits of Functions

## You will learn

- Unspecified quantities
- Limit of a function at a point.

## Key terms

- Unspecified quantity
- Undefined
- Value of a function
- Limit of a function

## Materials

- Scientific calculator.
- Graphical programs for computer

The concept of the function limit at a point is one of the basic concepts in calculus. This concept basically depends on the behavior of the function at all of its definition points. To study such a behavior, we should identify the types of quantities in the set of real number.



## Think and discuss

Find the sum of the following operation, if possible:

1  $3 \times 5$

2  $28 \div 4$

3  $4 - 9$

4  $7 \div 0$

5  $0 \div 0$

6  $\infty + 3$

7  $\infty \div \infty$

8  $\infty - \infty$



## Remember

$\infty$  is a symbol that indicates an unspecified quantity greater than any real number that can be imagined.

## Unspecified Quantities



## Learn

In **Think and discuss** above, we found that some sums of the operations are identified completely such as 1, 2, 3 while others can't be identified such as other operation.

**Notice:**  $7 \div 0$  is undefined where the division by zero does not make a sense.

Now, you can't determine the sum of the operation  $0 \div 0$  where there are an infinite numbers of numbers if multiplied by zero, then the product will be zero. Therefore,  $\frac{0}{0}$  is unspecified quantity:

$\frac{\infty}{\infty}$ ,  $\infty - \infty$  and  $0 \times \infty$  are from the unspecified quantities. (why?)



## Add to your information

Mathematical operations are performed on the set of real numbers and the two symbols  $\infty$  and  $-\infty$  as follows:

1  $\infty + a = \infty$

2  $-\infty + a = -\infty$

3  $\frac{a}{\infty} = \frac{a}{-\infty} = 0$

4  $\infty \times a = \begin{cases} -\infty, & \text{if } a < 0 \\ \infty, & \text{if } a > 0 \end{cases}$

5  $-\infty \times a = \begin{cases} -\infty, & \text{if } a > 0 \\ \infty, & \text{if } a < 0 \end{cases}$

### Example

1 Find the sums of the following operations (if possible):

a  $4 + \infty$

b  $3 - \infty$

c  $0 \div 3$

d  $-5 \div 0$

e  $\infty + \infty$

f  $0 \div 0$

g  $5 \times \infty$

h  $-6 \times -\infty$

### Solution

a  $\infty$

b  $-\infty$

c 0

d undefined

e 0

f unspecified quantity

g  $\infty$

h  $\infty$

### Try to solve

1 Find the sums of the following operations (if possible):

a  $0 \div (-2)$

b  $7 \div 0$

c  $9 \div \infty$

d  $\infty \times 0$

e  $(-7) \times \infty$

f  $(-\infty) + 12$

g  $\infty + \infty$

h  $\infty \div \infty$

### The limit of function at a point :



### Activity

In the following figure: the graphical line of the function  $f$  defined in  $\mathbb{R}$  according to the rule  $f(x) = 2x + 1$ . Complete the following tables, then answer the following questions:

$x$	$f(x)$
2,1	5,2
2,01	5,02
2,001	5,002
2,0001	5,0002
.....	.....
↓	↓
2	5


$$x > 2$$

$x$  gets near to 2 in the right direction

$x$	$f(x)$
1,9	4,8
1,99	4,98
1,999	4,998
1,9999	4,9998
.....	.....
↓	↓
2	5

$$x < 2$$

$x$  gets near to 2 in the left direction



### Note:

- What is the value which  $f(x)$  gets near to when  $x$  gets near to 2 in the right direction?
- What is the value which  $f(x)$  gets near to when  $x$  gets near to 2 in the left direction?

When  $x$  gets near to the number (2) from right and left,  $f(x)$  gets near to the number (5). We express that mathematically as follows:  $\lim_{x \rightarrow 2} (2x + 1) = 5$

## Definition

If the value of the function  $f$  gets near to a unique value  $L$  when  $x$  gets near to a form left side and right side, then the limit of  $f(x)$  equals  $L$ . It is **written symbolically** as:  $\lim_{x \rightarrow a} f(x) = L$

1

**and read as:** Limit of  $f(x)$  when  $x$  gets near to a equals  $L$ .


## Example

- 2 If  $f(x) = \frac{x^2 - 4}{x - 2}$ , study the values of  $f(x)$  when  $x$  gets near to 2.

## Solution

$x$	$f(x)$
2,1	4,1
2,01	4,01
2,001	4,001
.....	.....
↓	↓
2	4
$x > 2$	

$x$	$f(x)$
1,9	3,9
1,99	3,99
1,999	3,999
.....	.....
↓	↓
2	4
$x < 2$	



From the graph and data shown in the table above, we find that  $f(x) \rightarrow 4$  when  $x \rightarrow 2$  from the right and left directions  $\therefore \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = 4$

## Notice:

- The hole in the graph illustrates a case of unspecified cases of  $\frac{0}{0}$  when  $x = 2$  (i.e. the function is undefined when  $x = 2$ )
- The presence of a limit to the function when  $x \rightarrow 2$  does not necessarily mean the function is defined when  $x = 2$ .

## P Try to solve

- 2 If  $f(x) = \frac{x^2 - 1}{x + 1}$ , then study the values of  $f(x)$  when  $x$  gets near to  $(-1)$

## Example

- 3 Find the  $\lim_{x \rightarrow 3} f(x)$  in each of the following figures:




fig (1)




fig (2)




fig (3)

**Solution**

Fig (1)  $\lim_{x \rightarrow 3} f(x) = 1$

Fig (2)  $\lim_{x \rightarrow 3} f(x) = 1$  (Notice that the function is not defined when  $x = 3$ )

Fig (3)  $\lim_{x \rightarrow 3} f(x)$  is not existed

**Try to solve**

- 3 Find the  $\lim_{x \rightarrow 1} f(x)$  in each of the following figures:




Fig (1)




Fig (2)




Fig (3)

**From previous examples, we conclude that:**

The presence of a limit to the function when  $x \rightarrow a$  does not necessarily mean the function is defined when  $x = a$  and vice versa if the function is defined when  $x = a$ , that does not necessarily mean the function has a limit when  $x = a$ .

**Verbal expression:** express by your manner the difference between the value of a function at a point and limit of that function at the same point.


**Exercises (3 - 1)**

**First: exercises on finding the limit graphically:**

- 1 From the graph, find:

a  $\lim_{x \rightarrow 0} f(x)$


b  $f(0)$



- 2 From the opposite graph, find the following if possible:


a  $\lim_{x \rightarrow 3} f(x)$

b  $f(3)$




3 From the opposite graph, find:

- a  $\lim_{x \rightarrow -2} f(x)$
- b  $f(-2)$
- c  $\lim_{x \rightarrow 0} f(x)$
- d  $f(0)$



4 The opposite graph illustrates the function  $f(x) = 2 - x^2$ . Find:


- a  $\lim_{x \rightarrow 0} (2 - x^2)$
- b  $f(0)$



$$f(x) = 2 - x^2$$

5 The opposite graph illustrates the function  $f(x) = \frac{x^2 - 4}{x + 2}$ . Find:


- a  $\lim_{x \rightarrow -2} f(x)$
- b  $f(-2)$



$$f(x) = \frac{x^2 - 4}{x + 2}$$

6 From the opposite graph, find:

- a  $f(0)$
- b  $\lim_{x \rightarrow 0} f(x)$
- c  $f(2)$
- d  $\lim_{x \rightarrow 2} f(x)$





**Second: Finding the limit of a function algebraically:**

- 7 Complete the following table and conclude  $\lim_{x \rightarrow 2} f(x)$  where  $f(x) = 5x + 4$

$x$	1.9	1.99	1.999	$\longrightarrow$	2	$\longleftarrow$	2.001	2.01	2.1
$f(x)$				$\longrightarrow$	?	$\longleftarrow$			

- 8 Complete the following table and conclude  $\lim_{x \rightarrow -1} (3x + 1)$

$x$	-0.9	-0.99	-0.999	$\longrightarrow$	-1	$\longleftarrow$	-1.001	-1.01	-1.1
$f(x)$				$\longrightarrow$	?	$\longleftarrow$			

- 9 Complete the following table and conclude  $\lim_{x \rightarrow -1} \frac{x^2 - 1}{x + 1}$

$x$	-0.9	-0.99	-0.999	$\longrightarrow$	-1	$\longleftarrow$	-1.001	-1.01	-1.1
$f(x)$				$\longrightarrow$	?	$\longleftarrow$			

- 10 Complete the following table and conclude  $\lim_{x \rightarrow 2} \frac{x - 2}{x^2 - 4}$

$x$	1.9	1.99	1.999	$\longrightarrow$	2	$\longleftarrow$	2.001	2.01	2.1
$f(x)$				$\longrightarrow$	?	$\longleftarrow$			

# Finding the Limit of a Function Algebraically

## You will learn

- Limit of a polynomial function.
- Some theories of limits.
- Using long division in finding the value of the function limit.
- Using theory of:  $\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$

## Key terms

- Limit of a function
- Polynomial function
- direct substitution
- synthetic division
- conjugate

## Materials

- Scientific calculators.
- Graphic program for Computer

In this lesson, we are going to identify some methods and theories to help us calculate the limit of a function at a point directly without a need for establishing tables and find the limit numerically or graph the curve to find the limit graphically.



### Activity

If  $f_1(x) = x^2 + 1$  and  $f_2(x) = 2x + 3$ , find:

1 -  $f_1(1)$  and  $\lim_{x \rightarrow 1} f_1(x)$  (what do you notice?)

2 -  $f_2(0)$  and  $\lim_{x \rightarrow 0} f_2(x)$  (what do you notice?)



### Learn

## Limit of a Polynomial Function

theory

➤ If  $f(x)$  is a polynomial function and  $a \in \mathbb{R}$

then:  $\lim_{x \rightarrow a} f(x) = f(a)$



### Example

1 Find the limit for each of the following functions:

a  $\lim_{x \rightarrow 2} (x^2 - 3x + 5)$

b  $\lim_{x \rightarrow 3} (-4)$



### Solution

a  $\lim_{x \rightarrow 2} (x^2 - 3x + 5)$

$= 4 - 6 + 5 = 3$  (direct substitution)

b  $\lim_{x \rightarrow 3} (-4) = -4$

notice that  $f(x) = -4$  is constant for all the values of  $x \in \mathbb{R}$



### Try to solve

1 Find the limit for each of the following functions:

a  $\lim_{x \rightarrow 1} (2x - 5)$

b  $\lim_{x \rightarrow -2} (3x^2 + x - 4)$

**Theory**

If  $\lim_{x \rightarrow a} f(x) = L$        $\lim_{x \rightarrow a} X(x) = m$   
 then:

1-  $\lim_{x \rightarrow a} K f(x) = K.L$

where  $K \in \mathbb{R}$

2-  $\lim_{x \rightarrow a} [f(x) \pm X(x)] = l \pm m$

3-  $\lim_{x \rightarrow a} f(x) \cdot X(x) = L.M$

4-  $\lim_{x \rightarrow a} \frac{f(x)}{X(x)} = \frac{L}{M}$  if  $M \neq 0$

2

5-  $\lim_{x \rightarrow a} (f(x))^n = L^n$

where  $L^n \in \mathbb{R}$

**Example**

2 Find each of the following limits:

a  $\lim_{x \rightarrow -1} \frac{3x+7}{x^2+2x-5}$

b  $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\tan x}{x}$

**Solution**

a  $\lim_{x \rightarrow -1} \frac{3x+7}{x^2+2x-5} = \frac{\lim_{x \rightarrow -1} (3x+7)}{\lim_{x \rightarrow -1} (x^2+2x-5)} = \frac{3 \times -1 + 7}{(-1)^2 + 2(-1) - 5} = \frac{4}{-6} = -\frac{2}{3}$

b  $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\tan x}{x} = \frac{\lim_{x \rightarrow \frac{\pi}{4}} \tan x}{\lim_{x \rightarrow \frac{\pi}{4}} x} = \frac{1}{\frac{\pi}{4}} = \frac{4}{\pi}$

**Try to solve**

2 Calculate the following limits:

a  $\lim_{x \rightarrow 2} \frac{x^2-3}{2x+1}$

b  $\lim_{x \rightarrow \pi} x \cos x$

**Theory**

➤ If  $f(x) = X(x)$        $x \in \mathbb{R} - \{a\}$

and  $\lim_{x \rightarrow a} X(x) = l$     then  $\lim_{x \rightarrow a} f(x) = l$

**Example**


3 Find:  $\lim_{x \rightarrow 1} \frac{x^3-1}{x-1}$

**Solution**

We notice that  $f(x) = \frac{x^3-1}{x-1}$  is unspecified when  $x = 1$

by factorizing and dividing by similar non-zero factors, then

$f(x)$  can be written as:



$$f(x) = \frac{\cancel{(x-1)}(x^2+x+1)}{\cancel{(x-1)}} = x^2+x+1$$


$$= k(x)$$

So far, we find that  $f(x) = k(x)$  for each  $x \neq 1$

Since  $\lim_{x \rightarrow 1} k(x) = 3$  (polynomial)

According to the previous theory, we conclude that  $\lim_{x \rightarrow 1} f(x) = 3$

$$\therefore \lim_{x \rightarrow 1} \frac{x^3-1}{x-1} = 3$$



**P Try to solve**

3 Find:  $\lim_{x \rightarrow -2} \frac{x^3+8}{x+2}$

**Example**

4 Find:  $\lim_{x \rightarrow 1} \frac{x^3-2x^2+1}{x^2+x-2}$

**Solution**

We notice that the function of the numerator  $f(x) = 0$  by substituting  $x = 1$  and the function of the denominator  $Q(x) = 0$  by substituting  $x = 1$ . This means that the factor  $(x - 1)$  is common in both the numerator and the denominator. Due to the difficulty of factorizing the function of the numerator into one of its factors  $(x - 1)$ , we use the long division in order to find the other factor of the expression  $x^3 - 2x^2 + 1$  as follows:

$$\begin{array}{r|l} x-1 & x^3 - 2x^2 + 1 \\ \hline x^2 - x - 1 & x^3 - x^2 \\ \hline & 0 - x^2 + 1 \\ & - x^2 + x \\ \hline & 0 - x + 1 \\ & - x + 1 \\ \hline & \text{zero} \end{array}$$

Therefore:

$$\lim_{x \rightarrow 1} \frac{(x-1)(x^2-x-1)}{(x-1)(x+2)} = \lim_{x \rightarrow 1} \frac{x^2-x-1}{x+2} = -\frac{1}{3}$$

**P Try to solve**

4 a  $\lim_{x \rightarrow 2} \frac{x^3-x^2-5x+6}{x-2}$

b  $\lim_{x \rightarrow -3} \frac{x^3-10x-3}{x^2+2x-3}$

**Guide to solve**

in the long division operation:

(1) order all terms of the dividend and the divisor ascendingly or descendingly in the same way.

(2) Divide the first expression of dividend by the first expression of the divisor and quotient.

(3) Multiply quotient by the divisor and subtract the product from the divisor to get the remainder.

(4) Continue the same way till terminate the division operation.

### Example

#### Using conjugate:

5 Find the following limits:

a  $\lim_{x \rightarrow 4} \frac{\sqrt{x-3} - 1}{x-4}$

b  $\lim_{x \rightarrow 5} \frac{x^2 - 5x}{\sqrt{x+4} - 3}$

#### Solution

**Notice that:**  $f(x) = \frac{\sqrt{x-3} - 1}{x-4}$  is unspecified when  $x = 4$

Therefore, we search for a method to get rid of the factor  $(x - 4)$  in both the numerator and the denominator.

$$\begin{aligned} \lim_{x \rightarrow 4} \frac{\sqrt{x-3} - 1}{x-4} &\times \frac{\sqrt{x-3} + 1}{\sqrt{x-3} + 1} = \lim_{x \rightarrow 4} \frac{x-3-1}{(x-4)(\sqrt{x-3} + 1)} \\ &= \lim_{x \rightarrow 4} \frac{x-4}{(x-4)(\sqrt{x-3} + 1)} \\ &= \lim_{x \rightarrow 4} \frac{1}{\sqrt{x-3} + 1} = \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \text{b } \lim_{x \rightarrow 5} \frac{x^2 - 5x}{\sqrt{x+4} - 3} &= \lim_{x \rightarrow 5} \frac{x^2 - 5x}{\sqrt{x+4} - 3} \times \frac{\sqrt{x+4} + 3}{\sqrt{x+4} + 3} \\ &= \lim_{x \rightarrow 5} \frac{x(x-5)(\sqrt{x+4} + 3)}{x+4-9} = \lim_{x \rightarrow 5} \frac{x(x-5)(\sqrt{x+4} + 3)}{(x-5)} \\ &= \lim_{x \rightarrow 5} x(\sqrt{x+4} + 3) = 5(3+3) = 30 \end{aligned}$$

#### Try to solve

5 Find the following limits:

a  $\lim_{x \rightarrow 5} \frac{\sqrt{x-1} - 2}{x-5}$

b  $\lim_{x \rightarrow -1} \frac{x+1}{\sqrt{x+5} - 2}$

#### Theory

$$4 \quad \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = n a^{n-1}$$

### Example

6  $\lim_{x \rightarrow 1} \frac{x^{19} - 1}{x - 1} = \lim_{x \rightarrow 1} \frac{x^{19} - 1^{19}}{x - 1} = 19 \times 1^{18} = 19$

#### Corollaries

#### Corollaries on the theory:

1-  $\lim_{x \rightarrow 0} \frac{(x+a)^n - a^n}{x} = n a^{n-1}$

2-  $\lim_{x \rightarrow a} \frac{x^n - a^n}{x^m - a^m} = \frac{n}{m} a^{n-m}$

 **Example**

7 Find:

$$\text{a } \lim_{x \rightarrow 0} \frac{(x+5)^4 - 625}{x}$$

$$\text{b } \lim_{x \rightarrow 2} \frac{x^5 - 32}{x^2 - 4}$$

$$\text{c } \lim_{x \rightarrow 0} \frac{(x+1)^{11} - 1}{x}$$

$$\text{d } \lim_{x \rightarrow 2} \frac{(x-4)^5 + 32}{x-2}$$

 **Solution**

$$\text{a } = \lim_{x \rightarrow 0} \frac{(x+5)^4 - 5^4}{x} = 4 \times 5^3 = 500 \quad \text{b } = \lim_{x \rightarrow 2} \frac{x^5 - 2^5}{x^2 - 2^2} = \frac{5}{2} \times 2^3 = 20$$

$$\text{c } \lim_{x \rightarrow 0} \frac{(x+1)^{11} - 1^{11}}{x} = 11 \times 1^{11-1} = 11$$

$$\begin{aligned} \text{d } \lim_{x \rightarrow 2} \frac{(x-4)^5 + 32}{x-2} &= \lim_{x \rightarrow 2} \frac{(x-4)^5 - (-2)^5}{(x-4) - (-2)} \\ &= 5(-2)^4 = 80 \end{aligned}$$

 **Try to solve**

6 Find:

$$\text{a } \lim_{x \rightarrow -5} \frac{x^4 - 625}{x+5}$$

$$\text{b } \lim_{h \rightarrow 0} \frac{(h+3)^4 - 81}{h}$$

$$\text{c } \lim_{x \rightarrow 0} \frac{\sqrt[3]{x+1} - 1}{x}$$

 **Exercises (3 - 2)** 

Complete the following:

$$\text{1 } \lim_{x \rightarrow 2} (3x+1) = \dots$$

$$\text{2 } \lim_{x \rightarrow 1} \frac{x-1}{x+1} = \dots$$

$$\text{3 } \lim_{x \rightarrow 0} \frac{x^2 - x}{x} = \dots$$

$$\text{4 } \lim_{x \rightarrow 2} \frac{x^2 - 4}{x-2} = \dots$$

$$\text{5 } \lim_{x \rightarrow a} \frac{x^5 - a^5}{x-a} = \dots$$

$$\text{6 } \lim_{x \rightarrow 2} \frac{x^3 - 8}{x-2} = \dots$$

$$\text{7 } \lim_{x \rightarrow 1} \frac{x^2 + x - 2}{x-1} = \dots$$

$$\text{8 } \lim_{x \rightarrow 2} \frac{x^4 - 16}{x-2} = \dots$$

$$\text{9 } \lim_{x \rightarrow 2} \frac{x^5 - 32}{x^3 - 8} = \dots$$

$$\text{10 } \lim_{x \rightarrow 1} \left( \frac{x^2 - 1}{x-1} \right)^5 = \dots$$

$$11 \quad \lim_{x \rightarrow 2} \frac{x^2 - x - 2}{2x - 4} = \dots\dots$$

$$12 \quad \lim_{x \rightarrow -1} \frac{x^2 + 1}{x^3 + 1} = \dots\dots$$

Choose the right answer:

$$13 \quad \lim_{x \rightarrow 0} \frac{x^2 - 1}{x} \text{ equals:}$$

a 0

b 1

c 2

d has no limit

$$14 \quad \lim_{x \rightarrow -1} \frac{x^2 + x}{x + 1} \text{ equals:}$$

a -1

b 0

c 1

d 3

$$15 \quad \lim_{x \rightarrow 2} \frac{2x^2 - 8}{x - 2} \text{ equals:}$$

a 2

b 4

c 6

d 8

$$16 \quad \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sin x}{x} \text{ equals:}$$

a 1

b  $\frac{\pi}{2}$

c  $\frac{2}{\pi}$

d has no limit

$$17 \quad \lim_{x \rightarrow \frac{\pi}{4}} \frac{\tan x}{x} \text{ equals:}$$

a  $\frac{\pi}{2}$

b 1

c  $\frac{4}{\pi}$

d has no limit

Find the value for each of the following limits (if found)

$$18 \quad \lim_{x \rightarrow 3} (x^2 - 3x + 2)$$

$$19 \quad \lim_{x \rightarrow -2} \frac{x^2 + 1}{x - 3}$$

$$20 \quad \lim_{x \rightarrow \frac{\pi}{2}} (2x - \sin x)$$

$$21 \quad \lim_{x \rightarrow \pi} \frac{\cos 2x}{x}$$

$$22 \quad \lim_{x \rightarrow -1} \frac{x + 1}{x^3 + 1}$$

$$23 \quad \lim_{x \rightarrow 9} \frac{9 - x}{x^2 - 81}$$

$$24 \quad \lim_{x \rightarrow 4} \frac{x^2 + 4}{x - 4}$$

$$25 \quad \lim_{x \rightarrow -1} \frac{x^2 - 1}{x^2 + x}$$

$$26 \quad \lim_{x \rightarrow 4} \frac{4x^2 - 64}{x - 4}$$

$$28 \quad \lim_{x \rightarrow -1} \frac{x^2 - x - 2}{x + 1}$$

$$30 \quad \lim_{x \rightarrow -2} \frac{x^3 + 8}{x^2 - 4}$$

$$32 \quad \lim_{x \rightarrow 2} \frac{2x^2 - x - 6}{x^2 - x - 2}$$

$$34 \quad \lim_{x \rightarrow 0} \frac{(2x - 1)^2 - 1}{5x}$$

$$36 \quad \lim_{x \rightarrow 1} \left( \frac{x^2}{x+1} - \frac{3x+4}{x+1} \right)$$

$$38 \quad \lim_{x \rightarrow 1} \frac{x^3 + x^2 - 2}{x - 1}$$

$$40 \quad \lim_{x \rightarrow 1} \frac{x^3 + x - 2}{x^2 - 1}$$

$$42 \quad \lim_{x \rightarrow 3} \frac{\sqrt{4x - 3} - 3}{x - 3}$$

$$44 \quad \lim_{x \rightarrow 0} \frac{\sqrt{9x + 16} - 4}{x}$$

$$46 \quad \lim_{x \rightarrow 1} \frac{x^7 - 1}{x - 1}$$

$$48 \quad \lim_{x \rightarrow 4} \frac{x^3 - 64}{x - 4}$$

$$50 \quad \lim_{x \rightarrow 2} \frac{x^7 - 128}{2x - 4}$$

$$52 \quad \lim_{x \rightarrow 4} \frac{2x^3 - 128}{x^2 - 16}$$

$$54 \quad \lim_{x \rightarrow 0} \frac{(x+1)^9 - 1}{x}$$

$$56 \quad \lim_{x \rightarrow 1} \frac{(x+2)^4 - 81}{x - 1}$$

$$27 \quad \lim_{x \rightarrow 5} \frac{x^2 - 25x}{x - 5}$$

$$29 \quad \lim_{x \rightarrow -1} \frac{5x^2 + 5}{3x^2 - 3}$$

$$31 \quad \lim_{x \rightarrow \frac{3}{2}} \frac{2x^2 - x - 3}{4x^2 - 9}$$

$$33 \quad \lim_{x \rightarrow -3} \frac{2x^2 + 5x - 3}{x^2 + x - 6}$$

$$35 \quad \lim_{x \rightarrow 0} \frac{(x+1)^3 - 1}{x}$$

$$37 \quad \lim_{x \rightarrow 1} \frac{x^3 - x^2 + 2x - 2}{x - 1}$$

$$39 \quad \lim_{x \rightarrow 1} \frac{x^3 - 5x^2 - x}{x^4 + 2x}$$

$$41 \quad \lim_{x \rightarrow 1} \frac{\sqrt{x} - 1}{x - 1}$$

$$43 \quad \lim_{x \rightarrow 0} \frac{\sqrt{x+1} - 1}{x}$$

$$45 \quad \lim_{x \rightarrow 3} \frac{x - 3}{\sqrt{x} - \sqrt{3}}$$

$$47 \quad \lim_{x \rightarrow 2} \frac{x^4 - 16}{x - 2}$$

$$49 \quad \lim_{x \rightarrow 3} \frac{x^5 - 243}{x - 3}$$

$$51 \quad \lim_{x \rightarrow 2} \frac{x^6 - 64}{x^5 - 32}$$

$$53 \quad \lim_{x \rightarrow \frac{1}{2}} \frac{32x^5 - 1}{16x^4 - 1}$$

$$55 \quad \lim_{h \rightarrow 0} \frac{(3+h)^4 - 81}{6h}$$



# Limit of a Function at Infinity

In our life and practical applications, we enormously need to know the behavior of the function  $f(x)$  when  $x \longrightarrow \infty$ . The following activity shows that.




## Activity

Use a computer program to graph the function  $f$  where:  $f(x) = \frac{1}{x}$  and  $x > 0$ . What do you notice in the curve if the positive values of  $x$  increase to infinity? From the figure, we notice:

- The more the values of  $x$  increase and gets near to infinity, the more the values of  $f(x)$  get near to zero.

Therefore, we say that the limit of  $f(x)$  equals zero when  $x$  gets near to infinity.



## Learn

### Limit of a Function at Infinity

**Theory**

$$\lim_{x \rightarrow \infty} \frac{1}{x} = 0$$

**Corollary**

$$\lim_{x \rightarrow \infty} \frac{a}{x^n} = 0 \quad \{\text{where } n \in \mathbb{R}^+ \text{ and } a \text{ is a constant}\}$$

#### Basic rules:

- $\lim_{x \rightarrow \infty} c = c$ , where  $c$  is a constant
- if  $n$  is a positive number greater than 1, then  $\lim_{x \rightarrow \infty} x^n = \infty$

**Notice that the theory (2)** related to the limit of addition, subtraction, multiplication or division of two functions when  $x \longrightarrow \infty$  studied in the previous lesson is true when  $x \longrightarrow \infty$

#### You will learn

- Limit of a function at infinity.
- Find the limit of a function at infinity using algebraic solution.
- Find the limit of a function at infinity using graphical solution.



#### Key terms

- Limit of a function at infinity



#### Materials

- Scientific calculator.
- Pc graphical programs.

### Example

1 Find:

a  $\lim_{x \rightarrow \infty} \left(\frac{1}{x} + 3\right)$

b  $\lim_{x \rightarrow \infty} \left(4 - \frac{3}{x^2}\right)$

➤ Then check graphically using a graphical program.



### Solution

$$\begin{aligned} \text{a } \lim_{x \rightarrow \infty} \left(\frac{1}{x} + 3\right) &= \lim_{x \rightarrow \infty} \frac{1}{x} + \lim_{x \rightarrow \infty} 3 \\ &= 0 + 3 = 3 \end{aligned}$$

$$\therefore \lim_{x \rightarrow \infty} \left(\frac{1}{x} + 3\right) = 3$$

$$\begin{aligned} \text{b } \lim_{x \rightarrow \infty} \left(4 - \frac{3}{x^2}\right) &= \lim_{x \rightarrow \infty} 4 - \lim_{x \rightarrow \infty} \frac{3}{x^2} \\ &= 4 - 3 \lim_{x \rightarrow \infty} \frac{3}{x^2} = 4 - 3 \times 0 = 4 \end{aligned}$$

$$\therefore \lim_{x \rightarrow \infty} \left(4 - \frac{3}{x^2}\right) = 4$$



### Try to solve

1 Find:

a  $\lim_{x \rightarrow \infty} \left(\frac{5}{x} + 2\right)$

b  $\lim_{x \rightarrow \infty} \left(\frac{2}{x^2} + 5\right)$

### Example

2 Find:  $\lim_{x \rightarrow \infty} (x^3 + 4x - 5)$

**Solution**

$$\begin{aligned}\lim_{x \rightarrow \infty} (x^3 + 4x - 5) &= \lim_{x \rightarrow \infty} x^3 \left(1 + \frac{4}{x^2} - \frac{5}{x^3}\right), \text{ by taking off } x^3 \text{ a common factor.} \\ &= \lim_{x \rightarrow \infty} x^3 \times \lim_{x \rightarrow \infty} \left(1 + \frac{4}{x^2} - \frac{5}{x^3}\right) = \infty \times 1 = \infty\end{aligned}$$

**Try to solve**

2 Find each of the following limits:

**a**  $\lim_{x \rightarrow \infty} (x^3 + 7x^2 + 2)$

**b**  $\lim_{x \rightarrow \infty} (4 - 3x - x^3)$

**Example**

3 Find each of the following limits:

**a**  $\lim_{x \rightarrow \infty} \frac{2x - 3}{3x^2 + 1}$


**b**  $\lim_{x \rightarrow \infty} \frac{2x^2 - 3}{3x^2 + 1}$

**c**  $\lim_{x \rightarrow \infty} \frac{2x^3 - 3}{3x^2 + 1}$


**Solution**

 In all cases, divide both the numerator and the denominator by  $x^2$  (biggest power of the variable  $x$  in the denominator).


**a**  $\lim_{x \rightarrow \infty} \frac{2x - 3}{3x^2 + 1} = \frac{\lim_{x \rightarrow \infty} \left(\frac{2}{x} - \frac{3}{x^2}\right)}{\lim_{x \rightarrow \infty} \left(3 + \frac{1}{x^2}\right)} = \frac{0 - 0}{3 + 0} = 0$



**b**  $\lim_{x \rightarrow \infty} \frac{2x^2 - 3}{3x^2 + 1} = \frac{\lim_{x \rightarrow \infty} \left(\frac{2}{x} - \frac{3}{x^2}\right)}{\lim_{x \rightarrow \infty} \left(3 + \frac{1}{x^2}\right)} = \frac{2 - 0}{3 + 0} = \frac{2}{3}$



**c**  $\lim_{x \rightarrow \infty} \frac{2x^3 - 3}{3x^2 + 1} = \frac{\lim_{x \rightarrow \infty} \left(2x - \frac{3}{x^2}\right)}{\lim_{x \rightarrow \infty} \left(3 + \frac{1}{x^2}\right)} = \frac{\infty - 0}{3 + 0} = \infty$



**We deduce from this example that:** to find  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$  where both  $f(x)$  and  $g(x)$  are polynomial functions, then:

- The limit gives a real number not equal zero if the degree of the numerator equals the degree of the denominator.
- The limit equals zero if the degree of the numerator is less than the degree of the denominator.
- The limit gives  $(\infty \text{ or } -\infty)$  if the degree of the numerator is greater than the degree of the denominator.

**P Try to solve**

3 Find:

a  $\lim_{x \rightarrow \infty} \frac{5x^2 - 3x + 1}{2x}$

b  $\lim_{x \rightarrow \infty} \frac{4x^3 - 5x}{8x^4 + 3x^2 - 2}$

c  $\lim_{x \rightarrow \infty} \frac{-6x^2 + 1}{3x^2 + x - 2}$

**Example**

4 Find the following limits:

a  $\lim_{x \rightarrow \infty} \frac{x^3 - 2}{|x|^3 + 1}$

b  $\lim_{x \rightarrow \infty} (x - \sqrt{x^2 + 4})$

**Solution**

a  $\lim_{x \rightarrow \infty} \frac{x^3 - 2}{|x|^3 + 1}$

$\therefore x \rightarrow \infty$

$\therefore x > 0$  i.e.  $|x| = x$

$\therefore \lim_{x \rightarrow \infty} \frac{x^3 - 2}{x^3 + 1}$



$f(x) = \frac{x^3 - 2}{|x|^3 + 1}$

by dividing the numerator and the denominator by  $x^3$ 

$$= \frac{\lim_{x \rightarrow \infty} (1 - \frac{2}{x^3})}{\lim_{x \rightarrow \infty} (1 + \frac{1}{x^3})} = \frac{1 - 0}{1 + 0} = 1$$

b  $\lim_{x \rightarrow \infty} (x - \sqrt{x^2 + 4})$

$$= \lim_{x \rightarrow \infty} \frac{(x - \sqrt{x^2 + 4})}{1} \times \frac{(x + \sqrt{x^2 + 4})}{(x + \sqrt{x^2 + 4})}$$

$$= \lim_{x \rightarrow \infty} \frac{x^2 - x^2 - 4}{x + \sqrt{x^2 + 4}}$$

$$= \lim_{x \rightarrow \infty} \frac{-4}{x + \sqrt{x^2 + 4}}$$

$\therefore x \rightarrow \infty$

$\therefore x > 0 \rightarrow \sqrt{x^2} = |x| = x$  by dividing the numerator and the denominator by  $x = \sqrt{x^2}$

$$\therefore \lim_{x \rightarrow \infty} \frac{-4}{x + \sqrt{x^2 + 4}} = \frac{\lim_{x \rightarrow \infty} -\frac{4}{x}}{\lim_{x \rightarrow \infty} (1 + \sqrt{1 + \frac{4}{x^2}})} = \frac{0}{1 + 1} = 0$$



$f(x) = x + \sqrt{x^2 + 4}$

**P Try to solve**

4 Find the following limits:

a  $\lim_{x \rightarrow \infty} \frac{x - 3}{\sqrt{4x^2 + 25}}$

b  $\lim_{x \rightarrow \infty} (\sqrt{3x^2 + 5x} - \sqrt{3}x)$



## Exercises (3 - 3)



Complete the following:

- |   |   |
|---|---|
| ① $\lim_{x \rightarrow \infty} (1 + \frac{3}{x}) = \dots$                 | ② $\lim_{x \rightarrow \infty} (\frac{3}{x^2} - 2) = \dots$     |
| ③ $\lim_{x \rightarrow \infty} (-7) = \dots$                              | ④ $\lim_{x \rightarrow \infty} (x^2 - 3) = \dots$               |
| ⑤ $\lim_{x \rightarrow \infty} \frac{2x+1}{x} = \dots$                    | ⑥ $\lim_{x \rightarrow \infty} \frac{x^3-5}{x^2+1} = \dots$     |
| ⑦ $\lim_{x \rightarrow \infty} \frac{x^5+3}{x^3-5} = \dots$               | ⑧ $\lim_{x \rightarrow \infty} \frac{3x}{\sqrt{x^2-1}} = \dots$ |
| ⑨ $\lim_{x \rightarrow \infty} (3 - \frac{7}{x} + \frac{4}{x^2}) = \dots$ | ⑩ $\lim_{x \rightarrow \infty} (\sqrt{x^2+1} - x) = \dots$      |

Choose the correct answer:

- |   |      |                 |                 |            |
|---|------|-----------------|-----------------|------------|
| ⑪ $\lim_{x \rightarrow \infty} \frac{6x}{2x+3}$ equal:  | a 0  | b 2             | c 3             | d $\infty$ |
| ⑫ $\lim_{x \rightarrow \infty} \sqrt{\frac{4}{x}+1}$    | a 0  | b 1             | c 2             | d $\infty$ |
| ⑬ $\lim_{x \rightarrow \infty} \frac{x+3}{2-x^2}$       | a 0  | b $\frac{1}{2}$ | c $\frac{3}{2}$ | d $\infty$ |
| ⑭ $\lim_{x \rightarrow \infty} \frac{x^2+1}{2x-1}$      | a 0  | b $\frac{1}{2}$ | c 1             | d $\infty$ |
| ⑮ $\lim_{x \rightarrow \infty} \sqrt{\frac{1+x}{4x-1}}$ | a -1 | b $\frac{1}{4}$ | c $\frac{1}{2}$ | d 1        |

Find the limit of the function at infinity

- |  |  |   |
|--|--|---|
| ⑯ $\lim_{x \rightarrow \infty} \frac{3}{x^2}$          | ⑰ $\lim_{x \rightarrow \infty} (x^3 + 5x^2 + 1)$           | ⑱ $\lim_{x \rightarrow \infty} \frac{2-7x}{2+3x}$                 |
| ⑲ $\lim_{x \rightarrow \infty} \frac{x^2}{x+3}$        | ⑳ $\lim_{x \rightarrow \infty} \frac{4x^2}{x^2+3}$         | ㉑ $\lim_{x \rightarrow \infty} \frac{5-6x-3x^2}{2x^2+x+4}$        |
| ㉒ $\lim_{x \rightarrow \infty} \frac{2x-1}{x^2+4x+1}$  | ㉓ $\lim_{x \rightarrow \infty} \frac{x^3-2}{3x^2+4x-1}$    | ㉔ $\lim_{x \rightarrow \infty} \frac{2x^2-1}{4x^3-5x-1}$          |
| ㉕ $\lim_{x \rightarrow \infty} \frac{2x^2-6}{(x-1)^2}$ | ㉖ $\lim_{x \rightarrow \infty} (7 + \frac{2x^2}{(x+3)^2})$ | ㉗ $\lim_{x \rightarrow \infty} (\frac{1}{3x^2} - \frac{5x}{2+x})$ |

28.  $\lim_{x \rightarrow \infty} \left( \frac{x}{2x+1} + \frac{3x^2}{(x-3)^2} \right)$

29.  $\lim_{x \rightarrow \infty} \frac{-x}{\sqrt{4+x^2}}$

30.  $\lim_{x \rightarrow \infty} (\sqrt{4x^2 - 2x + 1} - 2x)$

31.  $\lim_{x \rightarrow \infty} (\sqrt{5x^2 + 4x + 7} - \sqrt{5x^2 + x + 3})$

32.  $\lim_{x \rightarrow \infty} \frac{x^2 + x - 1}{8x^2 - 3}$

33.  $\lim_{x \rightarrow \infty} \frac{4 - 3x^3}{\sqrt{x^6 + 9}}$

**Creative thinking**

A company produces post cards for a primary cost 5000 L.E and the cost of each card is half a pound. If the total cost is  $C = \frac{1}{2}x + 5000$  where  $x$  is the number of the produced cards,

**Find:**

- 1 The cost of the card when producing:
  - a 10000 cards
  - b 100000 cards
- 2 Find the cost of producing the card when the company produces an infinite number of cards.

**General Exercises**

For more exercises please visit the website of the Ministry of Education.


**Activity**

Use technology to find the limit of a function at a point (graphical calculator)

Use the graphical calculator to graph each of the following functions, then find the limit of each function at the given point:

**a**  $f(x) = x^3$  when  $x = 0$


**b**  $f(x) = \frac{x^3 - 1}{x - 1}$  when  $x = 1$

**c**  $f(x) = \frac{\sin^2 2x}{x^2}$  when  $x = 0$

**Solution**

- a** Use the graphical calculator to represent the curve of the function  $f(x) = x^3$ .


From graph,  $\lim_{x \rightarrow 0} f(x) = 0$ .



- b** Use the calculator to represent the curve of the function

$$f(x) = \left(\frac{x^3 - 1}{x - 1}\right) - 2$$


From graph,  $\lim_{x \rightarrow 1} f(x) = 1$  (note the hole at (1, 1))



- c** Use the calculator to represent the curve of the function

$$f(x) = \frac{\sin^2 2x}{x^2}$$

From the graph,  $\lim_{x \rightarrow 0} \frac{\sin^2 2x}{x^2} = 4$


**Enrichment Information**

Please visit the following links.



## Unit Summary

➤ Mathematical operations on the set of real numbers and the two symbols  $\infty$  and  $-\infty$  are done as follows:

①  $\infty + a = \infty$

②  $-\infty + a = -\infty$

③  $\frac{a}{\infty} = \frac{a}{-\infty} = \mathbf{zero}$

④  $\infty \times a = \begin{cases} -\infty, & \text{if } a < 0 \\ \infty, & \text{if } a > 0 \end{cases}$

⑤  $-\infty \times a = \begin{cases} -\infty, & \text{if } a > 0 \\ \infty, & \text{if } a < 0 \end{cases}$

➤ If the value of the function  $f$  gets near to a unique value  $L$  when  $x$  gets near to a form left and right sides, then the limit of  $f(x)$  equals  $L$ . It is written symbolically as:  $\lim_{x \rightarrow a} f(x) = L$  and read as: Limit of  $f(x)$  when  $x$  gets near to  $a$  equals  $L$ .

➤ The presence of a limit to the function when  $x \rightarrow a$  does not necessarily mean the function is defined when  $x = a$ , and vice versa if the function is defined when  $x = a$ , that does not necessarily mean the function has a limit when  $x = a$ .

➤ **If**  $\lim_{x \rightarrow a} f(x) = L$        $\lim_{x \rightarrow a} Q(x) = m$ , **then:**

①  $\lim_{x \rightarrow a} k f(x) = k.L$

where  $k \in \mathbb{R}$

②  $\lim_{x \rightarrow a} [f(x) \pm Q(x)] = l \pm m$

③  $\lim_{x \rightarrow a} f(x) \cdot Q(x) = L.M$

④  $\lim_{x \rightarrow a} \frac{f(x)}{Q(x)} = \frac{l}{m}$       if  $m \neq 0$

⑤  $\lim_{x \rightarrow a} (f(x))^n = l^n$       where  $l^n \in \mathbb{R}$

➤ limit of the function at infinity.

①  $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$

②  $\lim_{x \rightarrow \infty} \frac{a}{x^n} = 0$  {where  $n \in \mathbb{R}^+$  and  $a$  is a constant}

③  $\lim_{x \rightarrow \infty} c = c$ , where  $c$  is a constant if  $n$  is a positive integer, then  $\lim_{x \rightarrow \infty} x^n = \infty$

➤ To find  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$  where both  $f(x)$  and  $g(x)$  are polynomial functions, then:

① The limit gives a real number not equal zero if the degree of the numerator equals the degree of the denominator.

② The limit equals zero if the degree of the numerator is less than the degree of the denominator..

③ The limit gives  $(\infty$  or  $-\infty)$  if the degree of the numerator is greater than the degree of the denominator.

➤ to do a long division operation, we should consider the following:

① Order all expressions of the dividend and the divisor ascendingly and descendingly in the same way.

② Divide the first expression of dividend by the first expression of the divisor and quotient is written.

③ Multiply quotient by the divisor and subtract the product from the divisor to get the remainder.

④ Continue the same way till terminate the division operation.





## Cumulative Exam




- 1 Place each fraction from the following algebraic fraction in the simplest form :
- a  $\frac{x}{x^2 - x}$       b  $\frac{x+1}{x^2+2x+1}$       c  $\frac{x^2-25}{(x-5)^2}$       d  $\frac{x+3}{x^3-9x}$
- 2 Is  $k_1(x) = k_2(x)$ , if  $k_1(x) = \frac{2x}{2x+8}$ ,  $k_2(x) = \frac{x^2+4x}{x^2+8x+16}$ ? Explain.
- 3 If  $k_1(x) = \frac{4}{x+1}$ ,  $k_2(x) = \frac{3}{x-1}$ , find  $k(x) = k_1(x) + k_2(x)$  and show the domain of  $k$ .
- 4 Find the simplest form of the function  $f$  where  $f(x) = \frac{1}{x+1} + \frac{1}{x-1}$  and show its domain.
- 5 Find the simplest form of the function  $g$  where  $g(x) = \frac{x^2-1}{x^2} + \frac{x+5}{3x}$  and show its domain.
- 6 Write the symbolic expression for the following mathematical phrase:  
If  $f(x)$  gets near to  $L$  ( $L \in \mathbb{R}$ ) when  $x$  gets near to  $a$ , then  $L$  is defined as a limit of  $f(x)$  when  $x$  gets near to  $a$ .
- 7 If  $f(x) = \frac{x^2-1}{x-1}$ , study the values of  $f(x)$  when  $x$  gets near to 1.
- 8 If the function  $f$  where  $f(x) \begin{cases} x & \text{when } x < 2 \\ x+2 & \text{when } x \leq 2 \end{cases}$

Graph the curve of this function, then investigate the presence of  $\lim_{x \rightarrow 2} f(x)$

- 9 Give numerical examples to show the following:
- a Existence of limit of the function when  $x \rightarrow 1$  it does not necessarily mean that the function is defined at  $x = 1$ .
- b If the function is defined at  $x = 1$  it does not necessarily mean that there is an existence to the limit of the function.

- 10 From the opposite figure, find :

- a  $f(0)$       b  $\lim_{x \rightarrow 0} f(x)$
- c  $f(-2)$       d  $\lim_{x \rightarrow -2} f(x)$



- 11 Find the following limits, if found:

- a  $\lim_{x \rightarrow \infty} \frac{7x}{2x+5}$       b  $\lim_{x \rightarrow \infty} \frac{4x^2}{3-x}$       c  $\lim_{x \rightarrow \infty} \frac{|x|}{x}$       d  $\lim_{x \rightarrow \infty} \left( \frac{x^2+1-x}{x} \right)$
- e  $\lim_{x \rightarrow 2} \frac{x^2-5x+6}{x-2}$       f  $\lim_{x \rightarrow 1} \frac{\sqrt{x+3}-2}{x-1}$

## Unit 4

log

$e^x$   
ln

# Trigonometry

sin

cos

tan

## Unit introduction

Trigonometry ( a latin term ) is generally one of the mathematics branches and of geometry particularly where there is a relation among the side lengths of the triangle and the measurements of its angles in the form of trigonometrical functions( sine function , cosine function, tan function, ...). Ancient Egyptians had been the first to practically use the rules of trigonometry. They had used these rules to build their pyramids and temples. Our knowledge about trigonometry is related to Greek who had originated the rules and theories. Beroni had produced a proof to the area of the triangle in terms of its side lengths. Western civilization had known Euclidean geometry via Arabs. Arab people had a long history in trigonometry. They had used the six trigonometric ratios where Altabani had discovered the special relation of the spherical right-angled triangle and the rule of finding the rule of sunrise.

Trigonometry has a lot of life applications in calculating distances and angles used in constructing the buildings, playgrounds, roads and industry of engines and electrical and mechanical appliances. Furthermore, Trigonometry is used to calculate geographical and astronomical distances and the exploratory systems of satellites.

## Unit objectives

**By the end of this unit, student should be able to:**

- ◆ Identify the sine rule of any triangle that states in any triangle the side lengths are proportional to the sine of the opposite angles.
- ◆ Use the sine rule to find the side lengths of any triangle.
- ◆ Use the sine rule to find the measurements of the angles of any triangle.
- ◆ Deduce the relation between the sine rule of any triangle and radius length of its circumcircle.
- ◆ Deduce the cosine rule of any triangle.
- ◆ Use the cosine rule to find the unknown side length in the triangle.
- ◆ Use the cosine rule to find the unknown measurement of an angle in the triangle.
- ◆ Use sine and cosine rules of any triangle to solve this triangle:
- ◆ Use the calculator to solve various exercises and activities on sine and cosine rules of any triangle.

## Key terms

- Trigonometry
- Sine rule
- Cosine rule
- Acute angle
- Obtuse angle
- Right angle
- Shortest side
- Longest side
- Missing length
- UnKnown angle
- Smallest angle
- Largest angle
- The area of the triangle
- The side lengths of a Triangle
- Opposite angle

## Unit lessons


**Lesson (4-1):** Sine rule

**Lesson (4-2):** Cosine rule

## Materials

Scientific calculator

## Unit planning guide



**You will learn**

- ▶ The sine rule of any triangle.
- ▶ Using the sine rule to solve the triangle.
- ▶ Modeling and solving mathematical and life problems using the sine rule.
- ▶ The relation between the sine rule of any triangle, the radius length of the circumcircle of this triangle and solving problems on it.

**Key terms**

- ▶ Sine Rule
- ▶ Acute Angle
- ▶ Obtuse Angle
- ▶ Right Angle

**materials**

- ▶ Scientific calculator
- ▶ Graphical program

**Remember**

The inscribed angles which subtends the same arc in a circle are equal. The inscribed angle drawn in a semicircle is right angled.

**Preface**

You have learned how to solve the right-angled triangle. Now you are going to deal with triangles that don't have right angles to learn how to find the side lengths and the measurements of the angles of such triangles. You already know that each triangle is made up of six elements- three sides and three angles. If any three elements are given- a side length is to be involved at least, you can find the other three elements by using the sine and cosine rules. Here, we can say that we can solve the triangle.

**Learn****The sine rule**

The following figures represents three type of triangles.




Figure (1)

Figure (2)

Figure (3)

$$m(\angle A) = m(\angle D) \quad m(\angle A) = 180^\circ - m(\angle D) \quad m(\angle A) = 90^\circ$$

In figure (1) where  $\triangle ABC$  is an acute-angled triangle  $\sin A = \sin D = \frac{a}{2r}$

Similarly, we can deduce that:  $\sin B = \frac{b}{2r}$  and  $\sin C = \frac{c}{2r}$

In figure (2) where  $\triangle ABC$  is an obtuse-angled triangle at A

Notice:  $\sin A = \sin (180^\circ - D) = \sin D$

$$\sin (180^\circ - D) = \sin D$$

$$\sin D = \frac{a}{2r} \quad \therefore \sin A = \frac{a}{2r}$$

Similarly, we can deduce that :

$$\sin B = \frac{b}{2r} \quad \text{and} \quad \sin C = \frac{c}{2r}$$

«Check it with your teacher»

**Notice**

$a, b, c$  are symbols of the side lengths:  $\overline{BC}, \overline{AC}, \overline{AB}$  in  $\triangle ABC$  respectively.

**Now:** try to prove the same previous relation in  $\triangle ABC$  which is right - angled at A.

**In general** The sine rule in  $\triangle ABC$  states that:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2r \text{ where } r \text{ is the radius length of its circumcircle.}$$

i.e. In any triangle, the side lengths are proportional to the sines of the opposite angles of these sides



### Self learning

Prove the sine rule in different ways

**Use the sine rule to find the side lengths of any triangle.:**


### Example

- ① In the triangle ABC, If  $m(\angle A) = 75^\circ$ ,  $m(\angle B) = 34^\circ$  and  $a = 10.2$  cm, Find each of b and c to the nearest integer

### Solution

$$\therefore m(\angle A) + m(\angle B) + m(\angle C) = 180^\circ$$

$$\begin{aligned} \therefore m(\angle C) &= 180^\circ - (75^\circ + 34^\circ) \\ &= 71^\circ \end{aligned}$$



Use the sine rule to find b and c

$$\begin{aligned} \therefore \frac{a}{\sin A} &= \frac{b}{\sin B} = \frac{c}{\sin C} \\ b &= \frac{10.2 \times \sin 34^\circ}{\sin 75^\circ} \approx 6\text{cm} \end{aligned}$$

$$\therefore \frac{10.2}{\sin 75^\circ} = \frac{b}{\sin 34^\circ} = \frac{c}{\sin 71^\circ}$$

Use the calculator

1 0 . 2 × sin 3 4 ° ÷ sin 7 5 ° =

$$c = \frac{10.2 \times \sin 71^\circ}{\sin 75^\circ} \approx 10\text{cm}$$

Use the calculator

1 0 . 2 × sin 7 1 ° ÷ sin 7 5 ° =

### Try to solve

- ① In the triangle ABC, if  $m(\angle C) = 61^\circ$ ,  $m(\angle B) = 71^\circ$  and  $b = 91\text{cm}$ , find each of a and c.

### Finding the length of the longest side in the triangle

### Example

- ② Find the length of the longest side in the triangle ABC in which  $m(\angle A) = 49^\circ 11'$ ,  $m(\angle B) = 76^\circ 17'$  and  $c = 11.22\text{cm}$  to the nearest two decimals.

### Remember



The longest side of a triangle is the side opposite to the largest angle and vice versa the smallest angle in a triangle is the angle opposite to the shortest side.

**Solution**

$$\begin{aligned} \therefore m(\angle C) &= 180^\circ - [m(\angle A) + m(\angle B)] \\ &= 180^\circ - (49^\circ 11' + 76^\circ 17') = 54^\circ 32' \end{aligned}$$

The longest side is opposite to angle B. i.e., the required is to find b.

$$\begin{aligned} \therefore \frac{b}{\sin B} &= \frac{c}{\sin C} & \therefore \frac{b}{\sin 17^\circ 76'} &= \frac{11.22}{\sin 32^\circ 54'} \\ \therefore b &= \frac{11.22 \times \sin 17^\circ 76'}{\sin 54^\circ 32'} \approx 13.38 \text{ cm} \end{aligned}$$

**Try to solve**

- 2 Find the length of the shortest side in the triangle ABC in which  $m(\angle A) = 43^\circ$ ,  $m(\angle B) = 65^\circ$  and  $c = 8.4$  cm, to the nearest decimal.

**Solving the triangle using the sine rule**

Solving the triangle means that we find the measurements of its unknown elements if three elements of the six elements are known in a condition that one of those known elements is to be a side length at least because the triangle can not be solved if the measurements of its three angles are known only. The sine rule enables us to solve the triangle if the measurements of two angles and a side length are known (given).

**Solving the triangle if the measurements of two angles and a side length are given (known):**

Note: To solve the triangle ABC if the measurements of both angles B and C and the length of a are known, we follow the next steps:

- 1- We use the relation  $m(\angle A) + m(\angle B) + m(\angle C) = 180^\circ$  to find  $m(\angle A)$
- 2- We use the sine rule:  $\frac{a}{\sin A} = \frac{b}{\sin B}$  to find b
- 3- We use the sine rule:  $\frac{a}{\sin A} = \frac{c}{\sin C}$  to find c

The following examples illustrate that:

**Example**

- 3 Solve the triangle ABC in which  $m(\angle A) = 36^\circ$ ,  $m(\angle B) = 48^\circ$  and  $a = 8$  cm to the nearest three decimals.

**Solution**



Find  $m(\angle C)$  from the relation :

$$m(\angle C) = 180^\circ - (36^\circ + 48^\circ) = 96^\circ$$

We find b using the sine rule as follows:

$$\begin{aligned} \therefore \frac{a}{\sin A} &= \frac{b}{\sin B} & \therefore \frac{8}{\sin 36^\circ} &= \frac{b}{\sin 48^\circ} \\ \therefore b &= \frac{8 \times \sin 48^\circ}{\sin 36^\circ} & \therefore b &\approx 10.115 \text{ cm} \end{aligned}$$

Use the calculator as follows :  $8 \times \sin 48 \div \sin 36 =$



$$\begin{aligned} \therefore \frac{a}{\sin A} &= \frac{c}{\sin C} & \therefore \frac{8}{\sin 36^\circ} &= \frac{c}{\sin 96^\circ} \\ \therefore c &= \frac{8 \times \sin 96^\circ}{\sin 36^\circ} \approx 13.535 \text{ cm} \end{aligned}$$

Use the calculator as follows :

$$8 \times \sin 96 \div \sin 36 =$$

### Try to solve

- 3 Solve the triangle XYZ in which  $Y = 107.2 \text{ cm}$ ,  $m(\angle X) = 33^\circ 16'$  and  $m(\angle Z) = 44^\circ 19'$

### Geometrical applications

**The relation between the sine rule and the radius length of the circumcircle of this triangle.**

You have previously learned that:  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2r$  where  $r$  is the radius length of the circumcircle of this triangle.

### Example

- 4 ABC is a triangle in which  $a = 15 \text{ cm}$ ,  $m(\angle A) = 60^\circ$  and  $m(\angle B) = 45^\circ$ . Find  $C$  and the radius length of circumcircle of the triangle ABC, to the nearest integer.


### Solution

We find  $m(\angle C)$  as follows:

$$\begin{aligned} m(\angle C) &= 180^\circ - [60^\circ + 45^\circ] \\ &= 180^\circ - 105^\circ = 75^\circ \end{aligned}$$

We use the sine rule to find  $c$ :

$$\begin{aligned} \therefore \frac{c}{\sin C} &= \frac{a}{\sin A} & \therefore \frac{c}{\sin 75^\circ} &= \frac{15}{\sin 60^\circ} \\ c &= \frac{15 \times \sin 75^\circ}{\sin 60^\circ} \approx 17 \text{ cm} \end{aligned}$$



To find the radius length of the circumcircle of triangle ABC, we use the relation:

$$\frac{a}{\sin A} = 2r \quad \therefore \frac{15}{\sin 60^\circ} = 2r \quad \therefore 2r \times \sin 60^\circ = 15$$

$$\therefore r = \frac{15}{2 \sin 60^\circ} \approx 9 \text{ cm}$$

$$15 \div (2 \sin 60) =$$

### Try to solve

- 4 ABC is a triangle in which  $m(\angle A) = 64^\circ 23'$ ,  $m(\angle B) = 72^\circ 23'$  and  $c = 18 \text{ cm}$ . Find each of  $a$ ,  $b$  and the radius length of the circumcircle of triangle ABC.

### Example

- 5 ABC is a triangle inscribed in circle M with radius length 100 cm. If  $AB = AC = 182$  cm, find:
- Length of  $\overline{BC}$  to the nearest decimal.
  - The area of the triangle ABC to the nearest square centimetre.

### Solution

We find  $m(\angle B)$  as follows:

In  $\triangle ABC$  :

$$\frac{AC}{\sin B} = 2r \quad (\text{sine rule})$$

$$\frac{182}{\sin B} = 200 \quad \sin B = \frac{182}{200} = 0.91$$

$$\therefore m(\angle B) = 65^\circ 30' 19''$$

( $m(\angle B) = m(\angle C)$  because  $\triangle ABC$  is an isosceles triangle and both are acute angles)

We find  $m(\angle A)$

$$m(\angle A) = 180^\circ - 2 \times (65^\circ 30' 19'') \simeq 48^\circ 59' 22''$$

We find the length of  $\overline{BC}$  using the sine rule as follows:

$$\therefore \frac{BC}{\sin 48^\circ 59' 22''} = \frac{182}{\sin 65^\circ 30' 19''} \quad \therefore BC = \frac{182 \times \sin 48^\circ 59' 22''}{\sin 65^\circ 30' 19''} \simeq 150.9 \text{ cm}$$

$$1 \quad 8 \quad 2 \quad \times \quad \sin \quad 4 \quad 8 \quad \dots \quad 5 \quad 9 \quad \dots \quad 2 \quad 2 \quad \dots \quad ( \quad +$$

$$\sin \quad 6 \quad 5 \quad \dots \quad 3 \quad 0 \quad \dots \quad 1 \quad 9 \quad \dots \quad ( \quad =$$

$$\begin{aligned} \text{The area of the triangle } ABC &= \frac{1}{2} AB \times AC \sin A \\ &= \frac{1}{2} \times 182 \times 182 \sin 48^\circ 59' 22'' \simeq 12497 \text{ cm}^2 \end{aligned}$$

### Try to solve

- 5 ABC is a triangle inscribed in a circle with radius length 8.4 cm. If  $AB = AC = 10.3$  cm, Find:
- The length of the base  $\overline{BC}$
  - The area of the triangle ABC


### Daily life applications on the sine rule

The sine rule can be used to solve many applications. It could be done by drawing a triangle, then solving that triangle to find the required data.

### Rappel




The area of a triangle =  $\frac{1}{2}$  product of the lengths of any two sides  $\times$  sine the subtended angle.





### Example

- 6 **Sports:** the opposite figure represents three players from a football team during a match. Find the distance between the second player and the third player to the nearest feet.



**Hint**

The area of parallelogram ABCD =  $2A(\triangle ABC)$  and the area of  $\triangle ABC = \frac{1}{2} AB \times AC \sin BAC$

### Solution

$$m(\angle B) = 180^\circ - (70^\circ + 47^\circ) = 63^\circ$$

The distance between the second player and the third player is  $a$ .

$$\text{Then } \frac{a}{\sin 47^\circ} = \frac{92}{\sin 63^\circ} \quad \therefore a = \frac{92 \times \sin 47^\circ}{\sin 63^\circ} \approx 76 \text{ feet}$$

Use the calculator:

$$9 \quad 2 \quad \times \quad \sin \quad 4 \quad 7 \quad ( \quad \div \quad \sin \quad 6 \quad 3 \quad ( \quad =$$


The distance between the second player and the third player is approximately 76 feet

### Try to solve

- 6 Find the distance between the first player and the second player to the nearest feet.

### Example

- 7 **Geography:** In the following figure, there are three geographical positions forming a triangle. If the distance between position A and position B is 236 km, the measurement of the angle at position B is  $72^\circ$  and the measurement of the angle at position A is  $51^\circ$ , Find:



- The distance between position C and position B, to the nearest integer
- The area of land which positions A, B and C represent its vertices to the nearest square meter.

### Solution

- a We find  $m(\angle C)$  in  $\triangle ABC$  :  $m(\angle C) = 180^\circ - (51^\circ + 72^\circ) = 57^\circ$

We use the sine rule to find the length of  $\overline{BC}$  :

$$\therefore \frac{BC}{\sin A} = \frac{AB}{\sin C} \text{ (sine rule)} \quad \therefore \frac{BC}{\sin 51^\circ} = \frac{236}{\sin 57^\circ}$$

$$\text{then } BC = \frac{236 \times \sin 51^\circ}{\sin 57^\circ} = 218.6871 \approx 219 \text{ meters}$$


- b We find the surface area of the triangle ABC in terms of  $a$ ,  $b$  and  $m(\angle B)$ .

$$\begin{aligned} \therefore \text{The area of the triangle } ABC &= \frac{1}{2} a c \sin B \\ &= \frac{1}{2} \times 218.6871 \times 236 \times \sin 72^\circ \approx 24542 \text{ m}^2. \end{aligned}$$

**P Try to solve**

- 7 In the opposite figure, there are three geographical positions forming a triangle. If the distance between position X and position Y is 9 km, the measurement of the angle at position X is  $40^\circ$  and the measurement of the angle at position Z is  $105^\circ$ . Find :

- The distance between position X and position Z.
- The area of the triangle whose vertices are X, Y, Z.



Using the sine rule of any triangle to find the measurements of the angles (there are two possible solutions for an unknown angle).




**Activity 1**

Draw the triangle ABC in which  $b = 7$  cm,  $a = 5$  cm and  $m(\angle A) = 30^\circ$




**Materials:**

Paper - pencil - ruler - compasses - protractor.

- From point A, Draw  $\overrightarrow{AX}$
- From point A, use the protractor to draw a  $30^\circ$  angle with  $\overrightarrow{AX}$  then draw  $\overline{AC}$  of length 7 cm.
- Place the tip of the compasses at point C and open it for a distance of 5 cm to draw an arc intersecting  $\overrightarrow{AX}$  at point B. What do you notice?



We notice that the arc intersects  $\overrightarrow{AX}$  at two points. I.e. there are two graphs for the triangle ABC- one of them is a acute angled triangle and the other is an obtuse angled triangle.



- Compare the height of triangle (Z) drawn from point C  $\perp$   $\overrightarrow{AX}$  and the length of  $\overline{BC}$ . What do you notice?

**We notice that:**  $z = 3.5$  cm ,  $BC = 5$  cm and  $AC = 7$ cm i.e.  $z < a < b$

- Can you use the sine rule to find the measurements of the angles of the previous triangle? Explain. We investigate the possibility of solving the triangle ABC as follows:

We find the shortest dimension drawn from C on  $\overline{AB}$  , let it z.  $z = b \sin A$

$$\text{i.e. } z = 7 \sin 30 = 3 \frac{1}{2} \text{ cm}$$

Where  $\angle B$  is an acute angle and  $z < a < b$ , there are two values for angle B; one of them is an acute angle and the other is the complementary angle to it. We use the sine rule as follows:

$$\frac{b}{\sin B} = \frac{a}{\sin A} \quad \text{i.e.} \quad \frac{7}{\sin B} = \frac{5}{\sin 30} \quad \text{then} \quad \sin B = \frac{7 \times \sin 30^\circ}{5} = 0.7$$

Therefore  $m(\angle B) \simeq 44^\circ 25' 37''$

As a result, the other angle is (obtuse)  $\simeq 180^\circ - 44^\circ 25' 37'' \simeq 135^\circ 34' 23''$

### Application on the activity :


LMN is a triangle in which  $l = 12$  cm ,  $m = 15$  cm and  $m(\angle L) 40^\circ$ . Prove that the angle M has two values , then find them.

Use the Calculator to solve exercises and activities on the sine rule.



### Activity 2

The figure opposite represents three Egyptian cities forming a triangle. If the distance from Suiz to Cairo is 8cm and the measurement of the angle at Suiz is  $30^\circ$  and at Fayoum is  $40^\circ$ , find the distance from Cairo to Fayoum if each 1 cm in the figure represents 16,75 km in real to the nearest km.



- a) Can you find  $m(\angle A)$ ?

$$m(\angle A) = 180^\circ - (30^\circ + 40^\circ) = 110^\circ$$

$$= ( 0 4 + 0 3 ) - 0$$

$$8 1$$

- b) How can you find the real distance from Suiz to Cairo?

Length in reality = length in draw  $\div$  drawing scale

$$AB = 8 \div \frac{1}{16.75} = 134 \text{ km.}$$

$$8 \div 1 6 . 7 5 ( =$$

- c) How can you find the real distance from Cairo to Fayoum ?

We use the sine rule as follows :

$$\frac{b}{\sin B} = \frac{c}{\sin C}$$

### Remember



$$\text{Drawing scale} = \frac{\text{Length in drawing}}{\text{Length in reality}}$$

$$\text{Length in reality} = \frac{\text{Length in drawing}}{\text{drawing scale}}$$

$$\text{length in drawing} = \text{length in reality} \times \text{drawing scale}$$

i.e.  $\frac{b}{\sin 30^\circ} = \frac{134}{\sin 40^\circ}$  then  $= \frac{134 \times \sin 30^\circ}{\sin 40^\circ} \approx 104\text{km}$ .

1 3 4 × sin 3 0 ( + sin 4 0 ( =

- d Can you use the accurate length in the drawing to find the distance from Cairo to Fayoum?  
From the drawing of that activity, we find that :  $AC \approx 6.2$  cm  
Hence, the real length  $\approx 6.2 \div \frac{1}{16.75} \approx 104\text{km}$ .

**Drill on the activity:** In the activity above, use the sine rule to find the real distance from Suiz to Fayoum, then check your answer using the measurement.



### Exercises (4 - 1)



**Complete :**


- In any triangle , the side lengths are proportional to \_\_\_\_\_
- ABC is an equilateral triangle whose side length is  $10\sqrt{3}$  cm the diameter length of the circumcircle of this triangle is \_\_\_\_\_
- ABC is a triangle in which  $m(\angle A) = 60^\circ$ ,  $m(\angle C) = 40^\circ$  and  $c = 8.4$  cm, then  $a =$  \_\_\_\_\_ cm
- In the triangle ABC,  $\frac{2b}{\sin B} =$  \_\_\_\_\_ r
- The diameter length of the circumcircle of the acute triangle ABC is 20 cm and  $BC = 10\text{cm}$  then  $m(\angle A) =$  \_\_\_\_\_ °
- If the area of the equilateral triangle whose side length is 6 cm is \_\_\_\_\_

**Choose the correct answer .**


- The radius length of the circumcircle of the triangle ABC in which  $m(\angle A) = 30^\circ$  and  $a = 10$  cm is.....  
 a 10cm      b 20cm      c 5cm      d 40cm
- If the radius length of the circumcircle of the triangle ABC is 4 cm and  $m(\angle A) = 30^\circ$  , then the length of a is.....  
 a 4cm      b 2cm      c  $4\sqrt{3}$       d  $\frac{1}{16}$
- In the triangle ABC, the expression  $2r \sin A$  is equal to.....  
 a a      b b      c c      d  $A(\triangle ABC)$
- If r is the radius length of the circumcircle of the triangle XYZ , then  $\frac{y}{\sin Y}$  is equal to .....  
 a r      b 2 r      c  $\frac{1}{2}r$       d 4 r
- The triangle LMN in which  $m(\angle L) = 30^\circ$  and  $MN = 7$  cm, then the diameter length of its circumcircle is .....  
 a 7cm      b 3.5cm      c 14cm      d  $\frac{14}{\sqrt{3}}$

- 12 In the triangle XYZ, if  $3 \sin X = 4 \sin Y = 2 \sin Z$ , then  $x : y : z$  equal:  
 a 2 : 3 : 4      b 6 : 4 : 3      c 3 : 4 : 6      d 4 : 3 : 6
- 13 Use the sine rule to find the value of  $x$  to the nearest tenth.

a



b



Solve each triangle ABC by using the sine rule, if you know that:

- 14  $m(\angle A) = 75^\circ$ ,  $m(\angle B) = 34^\circ$ ,  $a = 10.2$ cm      15  $m(\angle A) = 19^\circ$ ,  $m(\angle C) = 105^\circ$ ,  $c = 11.1$ cm  
 16  $m(\angle A) = 116^\circ$ ,  $m(\angle C) = 18^\circ$ ,  $a = 17$ cm      17  $m(\angle A) = 36^\circ$ ,  $m(\angle B) = 77^\circ$ ,  $b = 2.5$ cm  
 18  $X(\angle A) = 49^\circ 11'$ ,  $m(\angle B) = 67^\circ 17'$ ,  $c = 11.22$ cm  
 19  $X(\angle B) = 115^\circ 4'$ ,  $m(\angle C) = 11^\circ 17'$ ,  $c = 516.2$ cm

Find the diameter length of the circumcircle of the triangle ABC in each of the following case :

- 20  $m(\angle A) = 75^\circ$ ,  $a = 21$ cm      21  $m(\angle B) = 50^\circ$ ,  $b = 90$ cm  
 22  $m(\angle C) = 102^\circ$ ,  $c = 11$ cm      23  $m(\angle A) = 70^\circ$ ,  $a = 8.5$ cm



### Activity

(24, 25; 26)




In each triangle ABC, find the measurements of the two angles B and C which satisfy the given conditions. Draw figures to help you determine whether there are two possible triangles or just one.

- 24  $m(\angle A) = 62^\circ$ ,  $a = 30$ cm,  $b = 32$ cm  
 25  $m(\angle B) = 48^\circ$ ,  $a = 93$  cm,  $b = 125$ cm  
 26  $m(\angle A) = 23.6^\circ$ ,  $a = 9.8$ cm,  $b = 17$ cm  
 27 In the triangle ABC,  $m(\angle A) = 67^\circ 22'$ ,  $m(\angle C) = 44^\circ 33'$  and  $b = 100$  cm, find the perimeter of the triangle ABC and its surface area.  
 28 In the triangle XYZ, If  $y = 68.4$ cm,  $m(\angle Y) = 100^\circ$  and  $X(\angle Z) = 40^\circ$ , find  $x$  and the radius length of the circumcircle of the triangle XYZ, then find the surface area of the triangle.  
 29 ABC is a triangle in which  $m(\angle A) = 22^\circ 37'$ ,  $m(\angle B) = 67^\circ 23'$ , and its perimeter is 30 cm. Find each of  $a$  and  $b$  to the nearest centimeter.

- 30 The perimeter of the triangle ABC is  $450 \text{ cm}^2$ ,  $m(\angle B) = 82^\circ$  and  $m(\angle C) = 56^\circ$ . Find the value of a.
- 31 ABCD is parallelogram in which  $AB = 18.6 \text{ cm}$ ,  $m(\angle CAB) = 36^\circ 22'$  and  $m(\angle DBA) = 44^\circ 38'$ , Find the diagonal length of  $\overline{AC}$  and the surface area of the parallelogram.
- 32 ABCD is a trapezium in which  $\overline{AD} \parallel \overline{BC}$ ,  $AD = 22.3 \text{ cm}$ ,  $m(\angle D) = 115^\circ$ ,  $m(\angle ACB) = 32^\circ 15'$  and  $m(\angle B) = 66^\circ$ , calculate the length of  $\overline{AC}$  and  $\overline{BC}$ .
- 33 ABCDE is a regular pentagon, whose side length is 18.36 cm. Find the diagonal length of  $\overline{AC}$ .
- 34  $\overline{AB}$  and  $\overline{AC}$  are two chords in a circle if their lengths are 43,5 cm and 52,1 cm and they are drawn in two different sides of the diameter  $\overline{AD}$  whose length is 100 cm, find:
- $m(\angle BAC)$
  - the length of  $\overline{BC}$
- 35 ABCD is a quadrilateral in which  $m(\angle BCD) = 85^\circ$ ,  $m(\angle CDA) = 87^\circ$ ,  $m(\angle BCA) = 36^\circ$ ,  $m(\angle BDA) = 55^\circ$  and  $CD = 100 \text{ cm}$ , find the length of  $\overline{BD}$  and  $\overline{AC}$ .
- 36 ABC is a triangle in which  $a = 58 \text{ cm}$ ,  $m(\angle B) = 38^\circ$  and  $m(\angle C) = 62^\circ$ , find the length of the perpendicular drawn from(A) on  $\overline{BC}$ .
- 37 A triangle - like piece of land A B C in which  $a = 90 \text{ meters}$ ,  $m(\angle B) = 53^\circ 8'$ ,  $m(\angle A) = 64^\circ 9'$ , find the perimeter and the area of this land.

### Creative thinking:

- 38 The distance between two light houses A and B is 20 km on one line from North to South. If a ship is located at position C where  $m(\angle ACB) = 33^\circ$  and  $m(\angle ABC) = 52^\circ$ , find the distance between the ship and each lighthouse.
- 39 **Climbing:** Adel and Karim stand in front of a rocky cliff to climb it and the distance between them is 8 meters as shown in the opposite figure. How high is the rocky cliff to the nearest tenth?
- 40 Ahmed and Salah stand in front of a minaret and the distance between them is 50 meters as shown in the opposite figure. How high is the minaret to the nearest tenth of meter?




## Cosine rule



## Think and discuss

Each of the following triangles is given the lengths of two sides ; 3cm and 4cm.



- From figure (1),  $m(\angle A)$  is a right angle. Find  $a$ .
- What are the possible values of " $a$ " if  $\angle A$  is an acute angle (figure 2)?
- What are the possible values of " $a$ " if  $\angle A$  is an obtuse angle (figure 3)?
- Can you solve the two triangles in figures (2) and (3) if  $(\angle A)$  is given using the sine rule? Explain.

Cosine rule helps us solve such triangles .



## Learn

## The cosine rule

In the figure opposite:  $\overline{CD} \perp \overline{AB}$

In  $\triangle BCD$ :

$$(BC)^2 = (CD)^2 + (BD)^2 \quad (\text{Pythagoras theory})$$


$$(BC)^2 = (CD)^2 + (AB - AD)^2 \quad \text{Expanding the brackets}$$

$$= (CD)^2 + (AD)^2 + (AB)^2 - 2AB \cdot AD$$

$$= (AC)^2 + (AB)^2 - 2AB \cdot AD$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

**Think:** Find the value of  $b^2$  and  $c^2$  in terms of  $a$ ,  $b$ ,  $c$  and the measurements of the angles of  $\triangle ABC$ .



## Notice



$$(AC)^2 = (AD)^2 + (CD)^2$$

$$AD = AC \cos A$$

## You will learn

- The cosine rule for any triangle.
- Using the cosine rule to solve the triangle.
- Modeling and solving daily life and mathematical problems using cosine rule.

## Key terms



- Cosine rule
- Acute angle
- Obtuse angle
- Right angle

## Materials



- Scientific calculator

The cosine rule states that : in any triangle ABC :

$$a^2 = b^2 + c^2 - 2bc \cos A, \quad b^2 = c^2 + a^2 - 2ca \cos B$$

$$, \quad c^2 = a^2 + b^2 - 2ab \cos C$$

### Critical thinking:

- 1 Prove the cosine rule when the triangle ABC is obtuse - angled.
- 2 Is the cosine rule true in case of the right - angled triangle? Explain.



### Activity 3

Log in internet or search in your school library for other proofs of the cosine rule, then discuss with your teacher.

### Finding the length of an unknown side in a triangle



### Example

- 1 XYZ is a triangle in which  $x = 24.3\text{cm}$ ,  $y = 22.8\text{cm}$  and  $m(\angle Z) = 42^\circ$ , find  $z$  to the nearest decimal.

### Solution

$$z^2 = x^2 + y^2 - 2x'y \cos Z$$

$$= (24.3)^2 + (22.8)^2 - 2 \times 24.3 \times 22.8 \cos 42^\circ \simeq 286.87$$

$$Z \simeq 16.9\text{ cm}$$

Use the calculator as follows:



### Try to solve

- 1 ABC is a triangle in which  $a = 72.8\text{ cm}$ ,  $b = 58.4\text{ cm}$  and  $m(\angle C) = 64.8^\circ$ , find  $c$  to the nearest decimal

### Finding the measurement of an angle if its three side lengths are given

You have previously learned that :

$$a^2 = b^2 + c^2 - 2bc \cos A \quad (\text{Cosine rule})$$

$$\text{i.e. } 2bc \cos A = b^2 + c^2 - a^2$$

$$\text{then } \cos A = \frac{b^2 + c^2 - a^2}{2bc} \quad (\text{dividing by } 2bc)$$

We can deduce that:

$$\cos B = \frac{c^2 + a^2 - b^2}{2ca} \quad \text{and} \quad \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$




Using the cosine rule of any triangle to find the measurement of an unknown angle in this triangle.

### Example

- 2 From the opposite figure, find  $m(\angle C)$

#### Solution

$$\begin{aligned}\cos C &= \frac{a^2 + b^2 - c^2}{2ab} && \text{(cosine rule)} \\ &= \frac{(5)^2 + (8)^2 - (12)^2}{2 \times 5 \times 8} && \text{(by substituting)} \\ &= \frac{-55}{80}\end{aligned}$$



The calculator can be used as follows:

$$\begin{array}{cccccccccccccccccccc} 5 & \chi^2 & + & 8 & \chi^2 & - & 12 & \chi^2 & + & ) & 2 & \times & 5 & \times & 8 & ( & = \\ \text{SHIFT} & \cos & = & - & \end{array}$$

We notice that the cosine of the angle is negative, so  $\angle C$  is obtuse and  $m(\angle C) \simeq 133^\circ 25' 57''$

### Try to solve

- 2 From the opposite figure, find  $m(\angle A)$

### Example

- 3 Find the measurement of the largest angle in the triangle LMN if it is known that  $l = 7.5$  cm,  $m = 12.5$  cm and  $n = 17.5$  cm, then prove that:

$$\cos N - 3\sqrt{3} \sin N + 5 = 0$$

#### Solution

The largest angle is the angle opposite to the longest side, so  $\angle N$  is the largest angle in the triangle.

$$\text{Then : } \cos N = \frac{l^2 + m^2 - n^2}{2lm} = \frac{(7.5)^2 + (12.5)^2 - (17.5)^2}{2 \times 7.5 \times 12.5} = -\frac{1}{2}$$


$$\therefore \cos N = -\frac{1}{2} \quad \therefore m(\angle N) = 120^\circ$$

$$\begin{array}{cccccccccccccccccccc} 7 & - & 5 & \chi^2 & + & 12 & - & 17 & \chi^2 & - & 7 & - & 5 & \chi^2 & = \\ + & ) & 2 & \times & 7 & - & 5 & \times & 12 & - & 17 & - & 5 & ( & = \\ \text{SHIFT} & \cos & \text{ANS} & ( & = & - & \end{array}$$

$$\begin{aligned}\text{The left side} &= \cos N - 3\sqrt{3} \sin N + 5 = \cos 120^\circ - 3\sqrt{3} \sin 120^\circ + 5 \\ &= -\frac{1}{2} - 3\sqrt{3} \times \frac{\sqrt{3}}{2} + 5 = 0 = \text{The right side.}\end{aligned}$$

### Try to solve

- 3 In the triangle ABC, if  $a = 12$  cm,  $b = 15$  cm and  $c = 18$  cm, prove that  $m(\angle C) = 2m(\angle A)$ .



### Remember

$$\begin{aligned}\cos 120^\circ &= \cos (180^\circ - 60^\circ) \\ &= -\cos 60^\circ = -\frac{1}{2} \\ \sin 120^\circ &= \sin (180^\circ - 60^\circ) \\ &= \sin 60^\circ = \frac{\sqrt{3}}{2}\end{aligned}$$

### Using the cosine rule in solving the triangle

The cosine rule allows us to solve the triangle in terms of the lengths of two sides and the measurement of the angle included. In this case, there is only one triangle.

### Solving the triangle in terms of the lengths of two sides and the measurement of the angle included :

#### Example

- 4 Solve the triangle ABC in which  $a = 11\text{cm}$ ,  $b = 5\text{cm}$  and  $m(\angle C) = 20^\circ$

#### Solution

$$\therefore c^2 = a^2 + b^2 - 2ab \cos C$$

$$\therefore c^2 = (11)^2 + (5)^2 - 2 \times 11 \times 5 \cos 20^\circ$$

$$\therefore c = \sqrt{(11)^2 + (5)^2 - 2 \times 11 \times 5 \cos 20^\circ}$$

$$\simeq 6.529\text{cm}$$

The calculator interface shows the following steps:  $\sqrt{\phantom{x}}$ , 11, 1,  $x^2$ , +, 5,  $x^2$ , -, 2,  $\times$ , 11, 1,  $\times$ , 5,  $\cos$ , 20, =.

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$= \frac{(5)^2 + (6.529)^2 - (11)^2}{2 \times 5 \times 6.529} \simeq -0.817$$

$$\therefore m(\angle A) \simeq 144.786^\circ$$

$$m(\angle B) = 180^\circ - [m(\angle A) + m(\angle C)]$$

$$= 180^\circ - [144.786^\circ + 20^\circ]$$


$$= 15.214^\circ$$

$$\therefore c = 6.529\text{cm}, m(\angle A) = 144^\circ 47' 96''$$

$$m(\angle B) \simeq 15^\circ 12' 50''$$

#### Try to solve

- 4 Solve the triangle ABC in which  $a = 24.6\text{cm}$ ,  $c = 14.2\text{cm}$  and  $m(\angle B) = 42^\circ 18'$



#### Remember

Solving the triangle means that we find the unknown elements. In this case, the required is to find  $c$ ,  $m(\angle A)$  and  $m(\angle B)$

#### Tip

To find the measurement of an angle in a triangle in terms of the lengths of two sides and the measurement of the angle included, it is better using the cosine rule instead of the sine rule. In case of using the sine rule, the sine of the acute angle or the obtuse angle is always positive because the sine is positive in the first and second quadrants. In case of using the cosine rule, if the angle is obtuse, its cosine is negative. If the angle is acute, its cosine is positive.

### Solving the triangle in terms of the lengths of its three sides

#### Example

- 5 Solve the triangle ABC in which  $a = 6$  cm,  $b = 8$  cm and  $c = 12$  cm.

#### Solution

Required is to find the measurements of the three angles of the triangle, then

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{(8)^2 + (12)^2 - (6)^2}{2 \times 8 \times 12} = \frac{43}{48}$$


$$\therefore m(\angle A) \simeq 26^\circ 23' 4''$$

$$\begin{array}{l} 8 \quad x^2 \quad + \quad 1 \quad 2 \quad x^2 \quad - \quad 6 \quad x^2 \quad = \quad + \quad ) \quad 2 \quad \times \\ 8 \quad \times \quad 1 \quad 2 \quad ( \quad = \quad \text{SHIFT} \quad \cos \quad \text{ANS} \quad ( \quad = \end{array}$$

$$\cos B = \frac{c^2 + a^2 - b^2}{2ca} = \frac{(12)^2 + (6)^2 - (8)^2}{2 \times 12 \times 6} = \frac{29}{36}$$

$$\therefore m(\angle B) \simeq 36^\circ 20' 10''$$

$$\begin{aligned} \therefore m(\angle C) &= 180^\circ - [26^\circ 23' 4'' + 36^\circ 20' 10''] \\ &= 117^\circ 16' 46'' \end{aligned}$$



#### Try to solve

- 5 Solve the triangle ABC in which  $a = 12.2$  cm,  $b = 18.4$  cm and  $c = 21.1$  cm

#### Writing in math :

Suppose you know the measurements of the three angles in a triangle. Can you use the sine rule or cosine rule to find the side length of this triangle? explain.

### Geometrical applications on the cosine rule


#### Example

- 6 ABC is a triangle in which  $a = 5$  cm,  $b = 2$  cm and  $c = 4$  cm. Bisect  $\overline{BC}$  at D, then draw  $\overline{AD}$  and find  $m(\angle C)$  and  $m(\angle CAD)$

#### Solution

In the triangle ABC

$$\begin{aligned} \cos C &= \frac{a^2 + b^2 - c^2}{2ab} \\ &= \frac{(5)^2 + (2)^2 - (4)^2}{2 \times 5 \times 2} = \frac{13}{20} \end{aligned}$$



$$\therefore m(\angle C) \simeq 49^\circ 27' 30''$$

$$\begin{array}{c} 5 \ x^2 + 2 \ x^2 - 4 \ x^2 = + ( 2 \times 5 \times 2 ) \\ = \text{SHIFT} \ \cos \ \text{ANS} \ = \end{array}$$

In the triangle ADC

$$\begin{aligned} (AD)^2 &= (DC)^2 + (AC)^2 - 2 DC \times AC \cos C \\ &= \left(\frac{5}{2}\right)^2 + (2)^2 - 2 \times \frac{5}{2} \times 2 \cos 30^\circ 27' 49'' \\ &\simeq 3.7499 \end{aligned}$$

$$\therefore AD \simeq 1.94 \text{ cm}$$

$$\begin{aligned} \therefore \cos(\angle CAD) &= \frac{(AC)^2 + (AD)^2 - (CD)^2}{2 \times AC \times AD} \\ &= \frac{(2)^2 + (1.94)^2 - (2.5)^2}{2 \times 2 \times 1.94} \simeq 0.1951 \end{aligned}$$

$$\therefore m(\angle CAD) \simeq 78^\circ 14' 14''$$

$$\begin{array}{c} 2 \ x^2 + 1 \ . \ 9 \ 4 \ x^2 - 2 \ . \ 5 \ x^2 = \\ ( 2 \times 2 \times 2 \times 1 \ . \ 9 \ 4 ) = \text{SHIFT} \ \cos \ \text{ANS} \ = \end{array}$$

### Example

- 7 **Geometry:** ABCD is a quadrilateral in which AB = 9 cm, BC = 5 cm, CD = 8 cm, DA = 9 cm and AC = 11 cm. Prove that the figure ABCD is a cyclic quadrilateral.

#### Solution

In the triangle ABC

$$\cos B = \frac{(9)^2 + (5)^2 - (11)^2}{2 \times 9 \times 5} = -\frac{1}{6}$$

In the triangle ADC

$$\cos E = \frac{(9)^2 + (8)^2 - (11)^2}{2 \times 9 \times 8} = \frac{1}{6}$$

Then  $\cos D = -\cos B$


$$\text{i.e. } m(\angle D) + m(\angle B) = 180^\circ$$

Where  $\angle D$  and  $\angle B$  are two opposite supplementary angles in the figure ABCD.

$\therefore$  The figure ABCD is a cyclic quadrilateral. (Q.E.D)

#### Try to solve

- 6 ABCD is a quadrilateral in which AB = 2.7 cm, AC = 7.2 cm, BC = 6.3 cm, CD = 4.5 cm and BD = 7.2 cm. Prove that the figure ABCD is a cyclic quadrilateral.






#### Remember



The cyclic quadrilateral is a figure whose four vertices belong to one circle

- the figure is cyclic quadrilateral if:
- There are two opposite supplementary angles.
- The measurement of the exterior angle at any vertex of its vertices equals the measurement of the interior angle opposite to the adjacent angle of this angle.
- There are two equal angles in measurement and drawn on one base and on one side of it.
- Its vertices are equidistant from a fixed point


**Discussion :** For each the following triangle, write the correct formula to the sine rule or cosine rule in order to find what is required (referred to in red). Use the given data referred to in blue only.



### Life applications on the cosine rule

#### Example

- 8 **Sports and tourism:** in the opposite figure, a tourist likes diving in order to watch the rare coral reefs and colorful fish. In a time of diving, the diver looked upward with an angle of  $20^\circ$ , he saw a squid of distance 3 meters. When he looked downward with an angle of  $40^\circ$ , he saw a red fish of a distance 4 meters. How far is it from the squid to the red fish?



#### Solution

It's clear from the drawing that we know the lengths of two sides in the triangle and the measurement of the included angle. Therefore, we can use the cosine rule as follows :

$$\begin{aligned} a^2 &= b^2 + c^2 - 2bc \cos A \\ &= (4)^2 + (3)^2 - 2 \times 4 \times 3 \cos 60^\circ \\ &= 13 \end{aligned}$$

$$\therefore a \simeq 3.6 \text{ m}$$

i.e. the distance from the squid to the red fish is approximately 3.6 meters.

#### Try to solve

- 7 **Sports:** Hany likes cycling. If he covers a distance of 6km from point A to point B, then covers another distance of 7 km from point B to point C where  $m(\angle ABC) = 79^\circ$ . How far is it from point A to point C to the nearest Km?

### Example

- 9 **Sports:** in a football match, the midfielder was at a distance of 20 meters from the right wing player. As the midfielder turned with an angle of measurement  $40^\circ$ , he saw the left wing player at a distance of 16 meters. What is the distance between the two wing players? to the nearest two decimals.


### Solution

Draw a figure to represent the problem as shown,

$$a^2 = b^2 + c^2 - 2bc \cos A$$


$$= (16)^2 + (20)^2 - 2 \times 16 \times 20 \cos 40^\circ \approx 12.87 \text{ meters}$$

The distance from the right wing to the left wing is approximately 12.87 meters




### Try to solve

- 8 **Games** in the bumper cars yard in an amusement park as shown in the figure. How far is it from car A to car B before they got collided?



### Example Measuring the distance indirectly

- 10 In the opposite figure, Shady wanted to measure the distance between the two points "A" and "B" on two different sides of a building from the location C which is at a distance of 33 meters from "A" and 48 meters from "B" as shown in the figure. If  $m(\angle C) = 54^\circ$ , find the distance AB to the nearest two decimal .



### Solution

In the triangle A B C the distance  $AB = c$


$$c^2 = a^2 + b^2 - 2 ab \cos C$$

$$= (48)^2 + (33)^2 - 2 \times 48 \times 33 \cos 54^\circ \approx 1530.8963$$

$$c \approx 39.13 \text{ meters}$$

### Try to solve

- 9 **Calculations of Land survey** Sanaa wanted to measure the distance from point "A" to point "B" on two different sides of a lake beach. She stands at location C which is at a distance of 258 meters from point "A" and 25.5 meters from point "B". If  $m(\angle C) = 78^\circ$ , find the length of  $\overline{AB}$  to the nearest two decimal.



### Exercises (4 - 2)





Complete the following:

- ① In any triangle XYZ,  $x^2 = y^2 + z^2 - \dots$ ,  $\cos X = \frac{y^2 + z^2 - \dots}{2yz}$
- ② A triangle whose side lengths are 13, 17 and 15 cm, then the measurement of its largest angle is \_\_\_\_\_
- ③ A triangle whose side lengths are 5,7 cm, 7,5 cm and 4,2 cm, then the measurement of its smallest angle is \_\_\_\_\_
- ④ A triangle ABC in which  $a = 10$  cm,  $b = 6$  cm and  $m(\angle C) = 60^\circ$ , then  $c =$  \_\_\_\_\_
- ⑤ In triangle LMN,  $m^2 + n^2 - \ell^2 =$  \_\_\_\_\_

Choose the correct answer:



- ⑥ The measurement of the largest angle in the triangle whose side lengths are 3, 5 and 7 is \_\_\_\_\_  
 a)  $150^\circ$       b)  $120^\circ$       c)  $60^\circ$       d)  $30^\circ$
- ⑦ In which triangle LMN, the expression  $\frac{\ell^2 + m^2 - n^2}{2\ell m}$  is equal to:  
 a)  $\sin L$       b)  $\cos M$       c)  $\cos N$       d)  $\sin N$
- ⑧ In triangle XYZ,  $y^2 + z^2 - x^2 = 2yz$  \_\_\_\_\_  
 a)  $\cos X$       b)  $\sin Z$       c)  $\cos Z$       d)  $\sin X$
- ⑨ In triangle ABC, if  $a : b : c = 3 : 2 : 2$ , then  $\cos a$  equals \_\_\_\_\_  
 a)  $\frac{1}{2}$       b)  $-\frac{1}{8}$       c)  $\frac{1}{2}$       d)  $\frac{3}{4}$

Use the cosine rule to find the value of  $x$  to the nearest tenth.




In triangle ABC if:

- 14  $a = 5$ ,  $b = 7$  and  $c = 8$ , prove that  $m(\angle B) = 60^\circ$
- 15  $a = 3$ ,  $b = 5$  and  $c = 7$ , prove that  $m(\angle C) = 120^\circ$
- 16  $a = 13$ ,  $b = 7$  and  $c = 13$ , find  $m(\angle C)$
- 17  $a = 13$ ,  $b = 8$  and  $c = 7$ , find  $m(\angle A)$
- 18  $a = 10$ ,  $b = 17$  and  $c = 21$ , find the measurement of the smallest angle.
- 19  $a = 5$ ,  $b = 6$  and  $c = 7$ , find the measurement of the largest angle.
- 20  $a = 17$ ,  $b = 11$  and  $m(\angle C) = 42^\circ$ , find  $c$  to the nearest two decimals.
- 21  $b = 16$  cm,  $c = 14$  cm,  $m(\angle A) = 72^\circ$ , find  $a$  to the nearest two decimals.
- 22 The triangle ABC in which  $a = 3$  cm,  $b = 5$  cm and  $c = \sqrt{19}$  cm, find :  
 a  $m(\angle C)$       b area of triangle ABC
- 23 ABC is a triangle in which  $a = 9$  cm,  $b = 15$  cm and  $c = 21$  cm, find the measurement of the largest angle of the triangle and prove that it satisfies the relation  $\cos C - 5\sqrt{3} \sin C + 8 = 0$
- 24 ABCD is a quadrilateral in which  $AB = 3$  cm,  $AC = 8$  cm,  $BC = 7$  cm,  $CD = 5$  cm and  $BD = 8$  cm, prove that the figure is a cyclic quadrilateral.
- 25 ABCD is a quadrilateral in which  $AB = 15$  cm,  $BC = 20$  cm,  $CD = 16$  cm,  $\overline{AC} = 25$  cm and  $m(\angle ACD) = 36^\circ 52'$ , find length of  $\overline{AD}$  to the nearest centimeter, then find the area of the quadrilateral ABCD.
- 26 ABCD is a parallelogram in which  $AB = 12$  cm,  $BC = 10$  cm and the diagonal length  $\overline{BD} = 14$  cm, find the diagonal length  $\overline{AC}$  to the nearest centimeter.
- 27 ABCD is quadrilateral in which  $BD = 78$  cm,  $CD = 96$  cm,  $m(\angle BCD) = 97^\circ$ ,  $m(\angle ABD) = 72^\circ$ , and  $m(\angle ADB) = 43^\circ$ , find  $\overline{AB}$ .
- 28 In triangle ABC,  $AB = 16$  cm,  $AC = 24$  cm and  $m(\angle A) = 80^\circ$ , find the length of  $\overline{BC}$ . If  $\overline{AD}$  bisects  $\angle A$  from inside, and intersects  $\overline{BC}$  at D, find the length of  $\overline{AD}$
- 29 **Sports :** A triangle - like race field whose side lengths are 1.2 km, 2 km and 1.8 km, find the measurement of each angle of its angles.
- 30 **Land survey :** A triangle - like piece of land whose side lengths are 300 m, 210 m and 140 m, use the cosine rule to find the area of the land to the nearest square meter.





- 31 **Sports** : Kareem rides his bicycle to cover a distance from point A to point B then to point C with speed of 28 km/h , then returns from point C to point A directly with speed of 35 km / h. How many minutes does he take back and forth to the nearest tenth?



- 32 **Writing in math** : Compare the two cases by which you can use the sine rule to solve the triangle and with the case you can use the cosine rule.
- 33 **Discover the error** : ABC is a triangle in which  $a = 5$  cm ,  $b = 10$  cm ,  $c = 7$  cm and  $m(\angle A) = 27.66^\circ$ , find  $m(\angle B)$ :

Ziad's solution

$$\begin{aligned} \therefore \frac{b}{\sin B} &= \frac{a}{\sin A} \\ \therefore \frac{10}{\sin B} &= \frac{7}{\sin 27.66^\circ} \\ \therefore \sin B &= \frac{10 \sin 27.66^\circ}{7} \approx 0.9488 \\ \therefore m(\angle B) &\approx 68.19^\circ \end{aligned}$$

Kareem's solution

$$\begin{aligned} \therefore \cos B &= \frac{c^2 + a^2 - b^2}{2ca} \\ \therefore \cos B &= \frac{(7)^2 + (5)^2 - (10)^2}{2 \times 7 \times 5} \approx -0.3714 \\ \therefore m(\angle B) &\approx 111.8^\circ \end{aligned}$$

### Critical thinking

- 34 If the two side lengths of a triangle are  $(\sqrt{10} + 2)$  and  $(\sqrt{10} - 2)$  and the included angle =  $60^\circ$ , find the third side length .
- 35 ABC is a triangle in which  $p - a = 8$  cm,  $p - b = 6$  cm and  $p - c = 4$  cm, find the measurement of the largest angle in the triangle where  $2p = a + b + c$ .
- 36 In triangle ABC , if  $p - a = 26$  cm,  $b = 28$  cm and  $p + a = 98$  cm where  $2p$  is the triangle perimeter, find the side lengths of the triangle then the measurement of the smallest angle.
- 37 If the ratio among the sines of the angles in a triangle is  $4 : 5 : 6$  , find the ratio among the cosines of these angles.
- 38 In triangle XYZ, if  $y^2 = (z - x)^2 + zx$  , prove that  $m(\angle Y) = 60^\circ$




### General Exercises

For more exercises please visit the website of the Ministry of Education.

## Unit summary


- 1 The triangle has six elements; three sides and three angles. .
- 2 Solving the triangle means that we find its unknown elements in terms of the given elements. In this unit, we have used the sine and cosine rules and the scientific calculator to solve the triangle and to solve daily life and geometrical applications.



- 3 **Sine rule :** in any triangle , the side lengths of the triangle are proportional to the sines of the angles opposite to them.

I.e. in any triangle ABC:  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

- This rule can be used to solve the triangle as the measurements of two angles and a side length are known or given.



- 4 In any triangle ABC, :

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2r$$

- Where r is the radius length of the circumcircle of triangle ABC.

- 5 **Cosine rule :**

- In triangle ABC :

- The cosine rule states that in any triangle ABC:

$$a^2 = b^2 + c^2 - 2bc \cos A \quad \text{then} \quad \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$b^2 = c^2 + a^2 - 2ca \cos B \quad \text{then} \quad \cos B = \frac{c^2 + a^2 - b^2}{2ca}$$

$$c^2 = a^2 + b^2 - 2ab \cos C \quad \text{then} \quad \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

- 6 **Using the cosine rule to solve the triangle :**

We can use the cosine rule for solving a triangle if known:

- Lengths of two sides and the measurement of the included angle.
- Lengths of the three sides.



- 7 **Area of the triangle :**  $\frac{1}{2}$  product of two adjacent sides by the sine of the angle included.

$$A(\triangle ABC) = \frac{1}{2} a b \sin C = \frac{1}{2} b c \sin A = \frac{1}{2} c a \sin B.$$

 **Enrichment Information**

Please visit the following link.




**Accumulative test**

**Choose the correct answer**


- 1 Without using the calculator, find the value of  $\cos 120^\circ$
- a  $-\frac{1}{2}$       b  $\frac{1}{2}$       c  $\frac{\sqrt{3}}{2}$       d  $\frac{2}{\sqrt{3}}$
- 2 Which of the following angles has negative sine and cosine?
- a  $75^\circ$       b  $135^\circ$       c  $265^\circ$       d  $330^\circ$
- 3 If  $\sin \theta = 46$ , then the measurement of angle  $\theta$  in degrees is:
- a 27.39      b -0.008      c 0.008      d 27.39
- 4 The relation links  $\tan E$  with  $\sec E$  is given in the form of :
- a  $\tan^2 E - 1 = \sec^2 E$       b  $\sec^2 E - 1 = \tan^2 E$       c  $\tan^2 E - \sec^2 E = 1$       d  $\tan^2 E + 1 = \sec^2 E$
- 5 Radius length of the circumcircle of triangle ABC in which  $m(\angle A) = 60^\circ$  and  $a = \sqrt{3}$  cm is:
- a 2cm      b  $\sqrt{3}$  cm      c  $2\sqrt{3}$       d  $\frac{\sqrt{3}}{2}$  cm
- 6 In any triangle LMN, the expression  $\frac{m^2 + n^2 - l^2}{2mn}$  equals
- a  $\cos L$       b  $\cos M$       c  $\sin L$       d  $\sin N$
- 7 In triangle ABC, b equals
- a  $\frac{c \sin B}{\sin C}$       b  $\frac{c \sin C}{\sin B}$       c  $\frac{c \sin B}{a}$       d  $\frac{c \sin C}{a}$
- 8 In triangle ABC, if  $a = 12$ ,  $b = 28$  and  $c = 20$ , then  $m(\angle B)$  equals
- a  $30^\circ$       b  $60^\circ$       c  $120^\circ$       d  $150^\circ$

**Short answer questions:**

- 9 Without using the calculator, find each of the following :
- a  $\cos 2\pi$       b  $\tan 135^\circ$       c  $\sin 330^\circ$       d  $\sec \frac{7\pi}{6}$
- 10 Find the accurate value for each of the following:
- a  $\sin (-300)^\circ$       b  $\sin 45^\circ \times \cos 210^\circ$       c  $\cos(\frac{7\pi}{6})$       d  $\sin \frac{3\pi}{2} \cos \frac{3\pi}{2}$
- 11 In triangle XYZ, if  $x = 10$  cm,  $m(\angle X) = 30^\circ$  and  $m(\angle Y) = 45^\circ$ , find  $y$ .
- 12 ABC is a triangle in which  $a = 4$  cm,  $b = 5$  cm and  $c = 6$  cm, find the measurement of the largest angle in the triangle then find its area.


- 13 In the figure opposite : use the given lengths in the triangle to check that:

a  $\sin^2 \theta + \cos^2 \theta = 1$    b  $\tan^2 \theta + 1 = \sec^2 \theta$




**Long answer questions:**


- 14 Solve the opposite triangle . Round the side length to the nearest tenth and the angle to the nearest degree.



(C)



(B)



(A)

- 15 XYZ is a triangle in which  $m(\angle X) = \frac{2}{3} m(\angle Y) = \frac{1}{2} m(\angle Z)$  and radius length of its circumcircle is 10 cm, find perimeter of the triangle XYZ.
- 16 Solve the triangle ABC in which  $a = 12$  cm,  $m(\angle C) = 66^\circ$  and  $c = 5$ cm. Round the length to the nearest centimeter and the angle to the nearest degree.
- 17 ABCD is a quadrilateral in which  $AB = 8$  cm,  $AD = 10$  cm,  $m(\angle A) = 82^\circ$  ,  $BC = 12$  cm, and  $m(\angle CBD) = 68^\circ$  , find the length of  $\overline{CD}$  to the nearest centimeter..

- 18 **History :** The greatest pyramid is the world's most controversial and imaginary monument. It is considered a real civilizational leap in Egypt's ancient history on that time, engineers had tried to build the pyramids frontage to be in the form of an equilateral triangle of a side length 230 meters. Find the height of the equilateral triangle to the nearest meter.



If you cant answer these questions, you can use the next table:

Question No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Back to	Prior knowledge	Prior knowledge	Prior knowledge	Prior knowledge	113	122	114	125	Prior knowledge	Prior knowledge	112	123	Prior knowledge	Prior knowledge	113	111	112	126

# General Exams


## Exam (1)

## Algebra

Answer the following questions:

**Question (1):** Choose the correct answer.

- ① If  $f : \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = x^3$ , then the figure which represents the function  $f$  is :



- ② If  $5^{x-3} = 4^{3-x}$ , then  $x =$

a  $\frac{5}{4}$                       b 3                      c  $\frac{4}{5}$                       d 0

- ③ The range of the function  $f$  where  $f(x) = |x|$  is

a  $[0, +\infty[$                       b  $]0, +\infty[$                       c  $] - \infty, 0]$                       d  $] - \infty, 0[$

- ④ If  $f(x) = 5^x$ , then  $f(-2) =$

a -2                      b 5                      c  $\frac{1}{25}$                       d  $\frac{1}{5}$

**Question (2):**

- ① If the function  $f$  where  $f(x) = \frac{1}{x}$ , find the range of the function  $f$ , the two coordinates of the symmetry point of the curve, then find the solution set of the equation  $f\left(\frac{1}{x}\right) = 4$

- ② Graph the curve of the function  $f$  where :

$$f(x) = \begin{cases} x^2 & \text{when } -5 \leq x < 2 \\ 6 - x & \text{when } 2 \leq x \leq 8 \end{cases}$$

From the graph, determine the range of the function and investigate its monotony.

**Question (3):**

- ① Graph the curve of the function  $f$  where  $f(x) = |x - 3|$ , deduce the range and monotony of the function and tell whether it is even, odd or otherwise.

2 Find the solution set for each in  $\mathbb{R}$  :

a  $|x - 3| \geq 5$

b  $|x - 3| = 0$

**Question (4):**

1 Find the solution set for each in  $\mathbb{R}$  :

a  $\log x = \log 3 + \log 10$

b  $9^x - 3 \times 3^x = 0$

2 Reduce:

a  $\frac{4^{2n+1} \times 2^{1-n}}{8^{n-2}}$

b  $\log_6 54 - \log_6 9$

**Question (5):**

1 Without using the calculator, find in the simplest form the value of :

$$\frac{1}{\log_2 30} + \frac{1}{\log_3 30} + \frac{1}{\log_5 30}$$

2 Tell whether each of the following two functions is odd or even :

a  $f(x) = x + \sin x$

b  $f(x) = x^3 - 2x^2$

## Exam (2)

## Algebra

Answer the following questions:

**Question (1): Choose the correct answer.**

1 The solution set of the inequality  $|x| - 1 > \text{zero}$  is:

a  $\mathbb{R} - [-1, 1]$

b  $[-1, 1]$

c  $\mathbb{R} - ]-1, 1[$

d  $[-1, 1]$

2 If  $4 = \log_2 x$ , then the equivalent exponential form is :

a  $x^2 = 4$

b  $x^4 = 2$

c  $x = 16$

d  $x = 8$


3 The domain of the function in the figure opposite is :

a  $[0, \infty[$

b  $]0, \infty[$

c  $[0, 1]$

d  $]0, 2[$



4 Which of the following functions represents an increasing exponential function on its domain  $\mathbb{R}$ :

a  $y = 3(1.05)^x$

b  $y = 3\left(\frac{1}{1.05}\right)^x$

c  $y = 3 + (0.5)^x$

d  $y = (0.05)^x$





- 2  $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1} =$
- a 0                      b 1                      c 2                      d 3
- 3 In  $\triangle ABC$ , if  $m(\angle A) = 30^\circ$  and  $a = 6$  cm, then  $\frac{b}{\sin B} =$  ...
- a 3                      b 6                      c  $\frac{1}{5}$                       d 12
- 4  $\lim_{x \rightarrow 1} \frac{x^5 - 1}{x - 1} =$  \_\_\_\_\_
- a 5                      b 1                      c 4                      d 20

**Question (2):**

1 Find:

a  $\lim_{x \rightarrow +\infty} \frac{5x^4 + 3x^2 - 6}{2x + x^4}$

b  $\lim_{x \rightarrow -2} \frac{x+2}{x-3}$

- 2 If ABC is
- $\triangle$
- in which :
- $\frac{1}{2} \sin A = \frac{1}{3} \sin B = \frac{1}{4} \sin C$
- , find its largest angle

**Question (3):**

1 Find :

a  $\lim_{x \rightarrow +\infty} \frac{4 - 3x^2}{\sqrt{x^4 + 5}}$

b  $\lim_{x \rightarrow 3} \frac{\sqrt{x+1} - 2}{x-3}$

- 2 Find the perimeter of
- $\triangle ABC$
- in which
- $a = 8$
- cm and
- $b = 6$
- cm and
- $m(\angle C) = 48^\circ$

**Question (4):**

1 Find:

a  $\lim_{x \rightarrow 3} \frac{x^2 - 6x + 9}{x - 3}$

b  $\lim_{x \rightarrow 2} \frac{2x^2 - 8}{x - 2}$

- 2 Find the diameter length of the circumcircle of
- $\triangle ABC$
- in the two following cases:

a  $m(\angle A) = 75^\circ$  ,  $a = 21$  cm

b  $m(\angle B) = 50^\circ$  ,  $m(\angle C) = 65^\circ$  ,  $c - b = 6$  cm

**Question (5):**

① Find the value of the following:

**a**  $\lim_{x \rightarrow 3} \frac{(x-6)^2 - 9}{x^2 - 9}$

**b**  $\lim_{x \rightarrow -1} \frac{2x^3 - x^2 - 2x + 1}{x^3 + 1}$

② ABC is a triangle in which  $m(\angle A) = 36^\circ$ ,  $m(\angle C) = 45^\circ$  and  $b = 9$  cm. Find the area of the circumcircle of the triangle.

**Exam (4)****Calculus and trigonometry**

Answer the following questions:

**Question (1): choose the correct answer.**

① In any triangle LMN :  $\frac{l}{\sin L}$  equals:

**a**  $\frac{m}{\sin N}$

**b**  $\frac{n}{\sin M}$

**c**  $\frac{m+n}{\sin N + \sin M}$

**d**  $3r$

②  $\lim_{x \rightarrow +\infty} \frac{\sqrt{4x^2 + 1}}{x - 2} =$

**a** 4

**b** 5

**c**  $\frac{5}{-2}$

**d** 2

③  $\lim_{x \rightarrow 0} (2x^2 + 3) =$

**a** 2

**b** 3

**c** 5

**d** 7

④ In  $\triangle ABC$ , if  $2 \sin A = 3 \sin B = 4 \sin C$ , then  $a : b : c$  equals \_\_\_\_\_

**a** 2 : 3 : 4

**b** 4 : 3 : 2

**c** 3 : 4 : 6

**d** 6 : 4 : 3

**Question (2):**

① Find:

**a**  $\lim_{x \rightarrow 2} \frac{x^5 - 32}{x - 2}$

**b**  $\lim_{x \rightarrow 1} \frac{(x-2)^4 - 1}{x-1}$

② ABCD is parallelogram in which  $AB = 7$  cm, the two diagonals  $\overline{AC}$  and  $\overline{BD}$  form two angles of measurements  $65^\circ$  and  $28^\circ$  with  $\overline{AB}$  respectively, find the length of  $\overline{BD}$  and  $\overline{AC}$

**Question (3):**

① Find :

**a**  $\lim_{x \rightarrow 3} \frac{x^3 - 27}{x^2 - 9}$

**b**  $\lim_{x \rightarrow \infty} \frac{4x^2 + 1}{x^2 - 2}$

② ABCD is a quadrilateral in which  $AB = 9$  cm,  $BC = 5$  cm,  $CD = 8$  cm,  $DA = 9$  cm and  $AC = 11$  cm, prove that ABCD is a cyclic quadrilateral .

**Question (4):****1 Find:**

**a**  $\lim_{x \rightarrow 1} \frac{x^2 + 5x - 6}{x^2 - 1}$

**b**  $\lim_{x \rightarrow 1} \frac{(x+1)^5 - 32}{x - 1}$

- 2** ABC is a triangle in which  $\cos A = \frac{2}{5}$ ,  $b = 2\frac{1}{2}$  and  $c = 2$ cm. Prove that the triangle is isosceles.

**Question (5):****1 Find:**

**a**  $\lim_{x \rightarrow 1} \frac{x^3 - 2x + 1}{x^2 - 1}$

**b**  $\lim_{x \rightarrow 1} \left( \frac{1}{x} + 3 \right)$

- 2** ABC is a triangle in which  $m(\angle B) = 35^\circ$ ,  $m(\angle C) = 70^\circ$ , and the radius length of the circumcircle of the triangle = 16cm, find the area and perimeter of triangle ABC to the nearest integer.

## المواصفات الفنية:

الرياضيات ص ٢ ع E

مقاس الكتاب:	$\frac{1}{8}$ (٨٢ × ٥٧) سم
طبع المتن:	٤ ألوان
طبع الغلاف:	٤ ألوان
ورق المتن:	٨٠ جم أبيض
ورق الغلاف:	٢٠٠ جم كوشيه
عدد الصفحات بالغلاف:	١٥٢ صفحة
رقم الكتاب:	١٥٣٨/١٠/١٥/١١/٢/٤٠
الكمية:	١٠٠١٠ نسخة

<http://elearning.moe.gov.eg>

---

**Kamal Fathalla Khedr Sons**

---